


# 2011 SAPMTCTE REPORT

## Early (4-8 weeks post-delivery) Population-level Effectiveness of WHO PMTCT Option A, South Africa, 2011



Medical Research Council, South Africa  
School of Public Health, University of the Western Cape,  
National Department of Health, South Africa  
Centers for Disease Control and Prevention/PEPFAR  
National Institute for Communicable Diseases/National Health Laboratory Service  
Wits Paediatrics HIV Diagnostics  
UNICEF



**2011 SAPMTCTE report**

**Early (4-8 weeks post-delivery)  
Population-level Effectiveness of WHO  
PMTCT Option A, South Africa, 2011**

***Principal Investigators***

Ameena Goga  
Thu-Ha Dinh  
Debra Jackson

***SAPMTCTE Study Group***

Yogan Pillay  
Gayle Sherman  
Adrian Puren  
Peter Barron  
Precious Robinson

Carl Lombard  
Selamawit Woldesenbet  
Vundli Ramokolo  
Tanya Doherty

Sanjana Bhardwaj  
Mary Mogoshoa

Nathan Shaffer  
Mickey Chopra

***Report Prepared by:***

Ameena Goga  
Debra Jackson  
Babalwa Dumalisile

***Contributors and Reviewers:***

Carl Lombard  
Adrian Puren  
Gayle Sherman  
Selamawit Woldesenbet  
Peter Barron  
Tanya Doherty  
Mary Mogoshoa  
Vundli Ramokolo

## **Copyright**

Copyright 2013. All material in this report may be reproduced and copied for non-commercial purposes: citation as to source, however, is required.

This report is disseminated by the South African Medical Research Council, National Department of Health South Africa and PEPFAR/US Centers for Disease Control & Prevention.

### **Suggested citation**

Goga AE, Dinh TH, Jackson DJ for the SAPMTCTE study group. Early (4-8 weeks post-delivery) Population-level Effectiveness of WHO PMTCT Option A, South Africa, 2011. South African Medical Research Council, National Department of Health of South Africa and PEPFAR/US Centers for Disease Control and Prevention. 2013.

ISBN: 978-1-920618-20-9

## CONTENTS

---

PRIMARY CONTACTS/PRINCIPAL INVESTIGATORS.....	vi
ACKNOWLEDGEMENTS.....	vii
ABBREVIATIONS AND ACRONYMS.....	viii
EXECUTIVE SUMMARY.....	ix
Introduction.....	ix
Aims and Objectives .....	ix
Methods.....	x
Results .....	x
Conclusions and Recommendations.....	xi
DEFINITIONS.....	xii
1. INTRODUCTION	1
2. METHODOLOGY	2
2.1 Survey Design and Justification .....	2
2.2 Study Population and Inclusion/Exclusion Criteria.....	3
2.3 Sampling .....	3
2.4 Data Collection Tools .....	4
2.5 Ethical Considerations .....	4
2.6 Data Collection Methods .....	5
2.7 Laboratory Methods.....	8
2.8 Quality Control of Field Work.....	9
2.9 Data Management.....	9
2.10 Data Analysis.....	10
3. RESULTS	11
3.1 Sample Realisation and Survey Profile .....	11
3.2 Sample Description and Characteristics .....	12
3.3 Infant HIV Infection Prevalence.....	19
3.4 National and Provincial Infant HIV Exposure and MTCT Rates.....	19
3.5 National PMTCT Programme Cascade.....	20
3.6 Demographic Characteristics, MTCT and the PMTCT Cascade by Province .....	24
3.7 Infant Feeding.....	58
4. DISCUSSION	61
4.1 Infant HIV Exposure .....	61
4.2 Mother-to-Child Transmission of HIV .....	61
4.3 PMTCT Cascade .....	63
4.4 Early Infant Diagnosis .....	64
4.5 Infant Feeding.....	64
5. STRENGTHS AND LIMITATIONS OF SAPMTCTE	65
Strengths.....	65
Limitations.....	65

6. CONCLUSION AND RECOMMENDATIONS	66
REFERENCES.....	67
SAMPLE SIZE CALCULATION BY PROVINCE.....	70
SAMPLING.....	71

## TABLES

Table 1 Data collection start and end dates in each province .....	5
Table 2 2010 SAPMTCTE desired and actual sample size by province .....	11
Table 3 Selected socio-demographic observations of 2011 SAPMTCTE .....	13
Table 4 Weighted Infant HIV infection prevalence nationally and by province .....	19
Table 5 Weighted infant HIV exposure and 4-8 week (early) MTCT of HIV by province for 2010 and 2011 .....	20
Table 6 HIV testing & results among pregnant women (weighted analysis) for 2010 and 2011 .....	21
Table 7 Access to the PMTCT programme in reported HIV-positive mothers (weighted analysis), 2010 and 2011 .....	22
Table 8 Baseline characteristics of Eastern Cape SAPMTCTE survey participants.....	24
Table 9 Baseline characteristics of Free State SAPMTCTE survey participants .....	27
Table 10 Baseline characteristics of Gauteng SAPMTCTE survey participants.....	31
Table 11 Baseline characteristics of KwaZulu-Natal SAPMTCTE survey participants .....	35
Table 12 Baseline characteristics of Limpopo SAPMTCTE survey participants.....	39
Table 13 Baseline characteristics of Mpumalanga SAPMTCTE survey participants .....	42
Table 14 Baseline characteristics of Northern Cape SAPMTCTE survey participants.....	46
Table 15 Baseline characteristics of North West SAPMTCTE survey participants.....	50
Table 16 Baseline characteristics of Western Cape SAPMTCTE survey participants.....	54
Table 17 Infant feeding practices amongst HIV exposed infants over the past 8 days by province ....	59

## FIGURES

Figure 1 Using ELISA at biomedical marker to identify HIV-exposed infants.....	2
Figure 2 Design phase and data collection flow diagram for the cell-phone data collection system ...	7
Figure 3 Example of SAPMTCTE Mobile Researcher web-based interface.....	7
Figure 4 2010 SAPMTCTE study profile.....	12
Figure 5 PMTCT service uptake (PMTCT cascade) in South Africa.....	23
Figure 6 PMTCT service uptake (PMTCT cascade) in the Eastern Cape .....	26
Figure 7 PMTCT service uptake (PMTCT cascade) in the Free State.....	30
Figure 8 PMTCT Service Uptake (PMTCT cascade) in Gauteng .....	34
Figure 9 PMTCT service uptake (PMTCT cascade) in KwaZulu-Natal.....	38
Figure 10 PMTCT service uptake (PMTCT cascade) in Limpopo .....	41

Figure 11 PMTCT service uptake (PMTCT cascade) in Mpumalanga .....	45
Figure 12 PMTCT service uptake (PMTCT cascade) in the Northern Cape .....	49
Figure 13 PMTCT service uptake (PMTCT cascade) in the North West Province .....	53
Figure 14 PMTCT service uptake (PMTCT cascade) in the Western Cape Province .....	57
Figure 15 Maternal antenatal HIV prevalence by province in South Africa.....	61
Figure 16 NHLS Early Infant Diagnosis PCR <2 months old 2008-2010 (from Sherman, 2010).....	62

---

Text Box 1: Eastern Cape infant HIV exposure and MTCT.....	26
Text Box 2: Free State infant HIV exposure and MTCT.....	29
Text Box 3: Gauteng infant HIV exposure and MTCT.....	33
Text Box 4: Kwa-Zulul Natal infant HIV exposure and MTCT.....	37
Text Box 5: Limpopo infant HIV exposure and MTCT.....	41
Text Box 6: Mpumalanga infant HIV exposure and MTCT.....	44
Text Box 7: Northern Cape infant HIV exposure and MTCT.....	48
Text Box 8: North West infant HIV exposure and MTCT.....	52
Text Box 9: Western Cape infant HIV exposure and MTCT.....	56

## PRIMARY CONTACTS/PRINCIPAL INVESTIGATORS

---

### **Ameena Goga, MD**

Paediatric Epidemiologist  
Medical Research Council, SA

*Address:* 1 Soutpansberg  
Road, Pretoria, 0001,

*Phone:* +2782 302 3168

*e-mail:*

Ameena.Goga@mrc.ac.za

### **Thu-Ha Dinh, MD, MS**

Medical Epidemiologist  
Centers for Disease Control and  
Prevention

*Address:* 1600 Clifton Rd  
Atlanta, 30333

*Phone:* +1 404 639 8618  
+2712 424 9000

*e-mail:* dvt1@cdc.gov;

dinhth@sa.cdc.gov

### **Debra Jackson, RN MPH DSc**

Professor (Extraordinary)  
School of Public Health  
Univ. of the Western Cape

*Address:* PBX17 Robert

Sobukwe Road, Bellville 7535

*Phone:* +2783 327 7331

*e-mail:*

debrajackson@mweb.co.za

## ACKNOWLEDGEMENTS

---

- CDC and PEPFAR for technical and financial support.
- UNICEF for financial support.
- National Department of Health and Global Fund.
- NICD/NHLS for conducting the validation ELISA testing on DBS samples; for providing consumables for the survey and for performing all the DBS ELISA and PCR testing. In particular we thank Ms Beverly Singh. Ushimta Patel and Ewalde Cutler.
- Ms Tsakani Mhlongo for training data collectors on infant blood drawing procedures.
- Provincial Departments of Health.
- District and facility managers who provided support for the SAPMTCT Evaluation.
- Mothers and their infants who participated in the survey.
- MRC survey supervisors and data collectors.
- WHO for protocol support.



## ABBREVIATIONS AND ACRONYMS

---

AIDS	Acquired Immunodeficiency Syndrome
ANC	Antenatal Care
ART	Antiretroviral therapy
ARV	Antiretroviral (drug)
CDC	Centers for Disease Control and Prevention
DBS	Dried Blood Spot
DHIS	District Health Information System
DHS	Demographic and Health Survey
DNA PCR	DNA-based Polymerase Chain Reaction Test
EBF	Exclusive Breast-Feeding
EID	Early Infant Diagnosis
ELISA	Enzyme-linked Immunosorbent Assay
HIV	Human Immunodeficiency Virus
HSRC	Human Sciences Research Council
HSRU	Health Systems Research Unit of the Medical Research Council
IMCI	Integrated Management of Childhood Illnesses
MCH	Maternal and Child Health
MDG	Millennium Development Goals
MPH	Masters in Public Health
MRC	Medical Research Council
MTCT	Mother-to-child transmission (of HIV)
NDOH	National Department of Health
NHLS	National Health Laboratory Service
NICD	National Institute for Communicable Diseases
NRF	National Research Foundation
NSP	National Strategic Plan, South Africa
PEPFAR	President's Emergency Plan For AIDS Relief
PITC	Provider-Initiated Testing and Counseling
PSU	Primary Sampling Unit
PMTCT	Prevention mother-to-child transmission of HIV
RtHC	Road to Health Chart
SA	South African
SAPMTCTE	South African Prevention of Mother-to-Child Transmission Evaluation
Sd-NVP	Single-dose Nevirapine
SoPH	School of Public Health, University of the Western Cape
UNICEF	United Nations Children's Fund
UWC	University of the Western Cape
WHO	World Health Organisation

## EXECUTIVE SUMMARY

---

### Introduction

The first national evaluation of the South African PMTCT programme was conducted between June and December 2010, during a period of transition between 2008 and 2010 guidelines to prevent HIV transmission from mother to child (PMTCT). The 2008 South African (SA) PMTCT guidelines offered maternal AZT prophylaxis from 28 weeks gestation and through labour and single dose nevirapine (sdNVP) or ART if  $CD4 \leq 250$  cells/ $\mu$ l or Stage IV disease. All infants received single dose nevirapine and 7 (or 28) days AZT.<sup>1</sup> The 2010 SA PMTCT guidelines offered maternal AZT prophylaxis from 14 weeks with FDC/TDF in labour and sdNVP or ART if  $CD4 \leq 350$  cells/ $\mu$ l or Stage III/IV disease.<sup>2</sup> All infants received six weeks nevirapine or for one week after breastfeeding stops.

The detailed report is available from <http://www.mrc.ac.za/healthsystems/reports.htm>. In summary a total of 10 820 eligible infants were enrolled; 10 735 interviews were conducted and 10 178 (94%) DBS were drawn and analysed.<sup>3</sup> The national weighted infant HIV-exposure prevalence was 32.0% (95% CI 30.7-33.3%). The national weighted MTCT rate measured at 4-8 weeks of infant age was 3.5% (95% CI 2.9-4.1%). Provincial MTCT ranged between 1.4% and 5.9%. Among mothers who reported being HIV negative, 4.1% delivered HIV-exposed infants. Of all participating mothers 98.8% (95% CI 98.5-99.0%) received an HIV test during pregnancy and of these 98.6% (95% CI 98.4-98.9) received their HIV test results. Of the reported HIV-positive mothers 78.3% had a CD4 cell count done during pregnancy and 91.8% received either maternal highly active antiretroviral therapy (ART) or mother/baby antiretroviral (ARV) prophylaxis. Amongst known HIV exposed infants 89% had received infant feeding counselling and only 35.1% intended to access early infant diagnosis services at the six weeks immunisation visit. Among all HIV exposed infants 20% were exclusively breastfeeding, 62% formula feeding and 18% mixed feeding in the 8 days prior to the interview.

The 2011 survey was the second national PMTCT evaluation to determine population-level PMTCT effectiveness, and the first national evaluation of WHO PMTCT Option A.

### Aims and Objectives

We aimed to conduct a national facility-based survey to monitor the effectiveness of the South African PMTCT programme. Our primary objective was to measure rates of early MTCT of HIV at six weeks postpartum. Our secondary objective was to estimate coverage of key PMTCT interventions and services (e.g., HIV testing, CD4 cell count testing, infant antiretroviral (ARV) prophylaxis, infant feeding counselling).

## Methods

The methods for the 2011 six weeks component was the same as in 2010.<sup>3</sup> The sampling frame and selected facilities were identical between 2010 and 2011, except for the 10 clinic replacements due to shifting of services or clinic closure for maintenance.

A cross-sectional facility-based survey was conducted at public primary health care/community health centres (PHC/CHC) offering immunisation services in all nine provinces. This methodology was chosen as uptake of six-week immunisation in South Africa, was >99%, according to the 2007 District Health Information System. The survey aimed to capture known and unknown HIV-exposed infants, as well as PMTCT participants and non-participants. A biomedical marker (HIV enzyme-linked Immunosorbent Assay (ELISA) tests to identify HIV antibodies) was used to identify HIV-exposed infants from infant dried blood spot (DBS) specimens. All DBS specimens reactive on ELISA testing were sent for DNA-based polymerase chain reaction tests (DNA PCR) to determine infant HIV infection status.

Infants aged 4-8 weeks attending PHC/CHC facilities for their six week immunisation were included. Hospitals and mobile clinics, very sick infants or infants aged <4 weeks or >8 weeks were excluded. The immunisation data from the 2007 District Health Information System (DHIS) were used to quantify the number of children that could be expected within facilities over a period of time and then stratify by size. Sample size was calculated so that valid national and provincial level estimates of MTCT could be ascertained. This resulted in between 34-79 facilities per province, 580 in total. Facilities were randomly selected within strata with probability proportional to size (3 strata). Caregiver/infant pairs were consecutively or randomly selected from facilities (depending on facility size). Interviews were conducted and infant DBS drawn after receiving consent from caregivers for study participation. Mothers and infants were referred into HIV care, as appropriate. Data were collected using low cost cell phones and interview data were uploaded real time into a web-based database console. Analysis was weighted for sample realisation and at provincial level proportional to the live birth distribution of South Africa.

## Results

In 2011 a total of 11 317 infants were screened from 580 facilities. Of these 601 did not meet eligibility criteria, 143 refused to participate and 158 had incomplete consent. 10 475 infants were deemed eligible and enrolled into the survey. 330 caregivers refused infant DBS and 39 had insufficient DBS that could not be analysed. Thus 10 106 DBS were analysed.

- The national weighted infant HIV-exposure prevalence was 32.0% (95% CI 30.7-33.3%).
- The national weighted MTCT rate measured at 4-8 weeks of infant age was 2.7% (2.1-3.2).
- The MTCT rate across provinces ranged from 1.98% to 6.06%..
- Among mothers who reported being HIV negative, 3.9% had HIV-exposed infants.
- Of all women participating 98.3 (95%CI 98.0-98.6) received an HIV test during pregnancy and of these 99.4 (95% CI 97.7-99.2) got their HIV test results.

- Amongst self-reported HIV-positive mothers:
  - 77.4% had a CD4 cell count done during pregnancy
  - 93.9% received either maternal highly active antiretroviral therapy (ART) or mother/baby antiretroviral (ARV) prophylaxis. 38.5% intended to access early infant diagnosis services at the six week immunisation visit and
  - 93.3% reported receiving infant feeding counselling.
- Amongst self-reported HIV-positive women, 35.5 (33.1-38.0%) were exclusively breastfeeding, 47.1% (44.8%-49.3%) avoided breastmilk and 17.4% (15.6-19.1%) were mixed feeding in the 8 days prior to the interview.
- Amongst self-reported HIV negative women 43.6% (41.6-45.7) were exclusively breastfeeding, 10.1% (9.3%-11.0%) avoided breastmilk and 46.2% (44.2-48.3%) were mixed feeding in the 8 days prior to the interview.

## Conclusions and Recommendations

1. The national PMTCT survey found a 2.7% national MTCT rate in pregnancy and intrapartum with a greater than 3-fold differential range of rates across the nine provinces (1.98% to 6.06%).
2. Maternal HIV acquisition since the last HIV test was potentially high at 3.9% and therefore repeat HIV testing at 32 weeks pregnancy, promotion of condom usage and couple testing is critical. Further data should be collected to assess the contribution of false negative rapid test results to maternal potential HIV acquisition. In addition, more work is required to improve the quality of rapid HIV testing in the field.
3. Uptake of PMTCT services is high, with more than 98% of women getting HIV tested during pregnancy and 93.9% of HIV-positive mothers receiving ARV treatment or prophylaxis. However CD4 (77.4%) testing and early infant diagnosis (EID) (38.5%) uptake are lower and represent on-going missed opportunities in the PMTCT programme.
4. Early infant HIV testing uptake is high if offered to all infants (98.28%) at six-week immunisation visits, indicating that EID strategies that routinely offer infant HIV testing only to known HIV-exposed infants should be reviewed.
5. Given the measured MTCT rate and continued reduction in MTCT compared to 2010 (3.5%) virtual elimination of paediatric HIV infection is possible with continued intensified effort. However, postnatal transmission after 6 weeks also needs to be examined to assess achievement of <5% MTCT at 18 months of infant age.
6. Exclusive breastfeeding was higher than previously reported, but was still low amongst self-reported HIV positive and negative women despite the adoption of the Tshwane Declaration of Support for Breastfeeding in August 2011.

## DEFINITIONS

---

Caregiver	The person who feeds and looks after the child most of the week. This includes parents, legal guardians, family members, nannies or friends who routinely feed, bath, change nappies, or in particular reference to this study, bring the child for routine health services.
The Consortium	Health Systems Research Unit (HSRU) of the Medical Research Council (MRC) and School of Public Health of the University of the Western Cape (SoPH, UWC).
Early (4-8 weeks) HIV transmission rate among HIV-exposed infants	Number of DNA PCR positive and ELISA positive infants divided by the number of ELISA positive infants at 4-8 weeks.
Health care personnel	Health care providers and health care workers.
Health care provider	Any person providing health services in terms of any law, including in terms of the: <ul style="list-style-type: none"><li>• Allied Health Professions Act, 1982 (Act No.63 of 1982),</li><li>• Health Professions Act, 1974 (Act No. 56 of 1974),</li><li>• Nursing Act, 2005 (Act No. 33 of 2005),</li><li>• Pharmacy Act, 1974 (Act No. 53 of 1974), and</li><li>• Dental Technicians Act, 1978 (Act No. 19 of 1979).</li></ul>
Health care worker	Any person who is involved in the provision of health services to a user, but is not a health care provider. This includes lay counselors and community caregivers.
HIV-exposed infant	An infant born to a known HIV-positive mother and/or having a positive HIV antibody test result using DBS ELISA. Infant HIV exposure prevalence serves as an indirect marker of maternal HIV prevalence.
HIV-infected infant	An HIV-exposed infant having a positive HIV DNA PCR result.
HIV-uninfected infant	An HIV-exposed infant having a negative HIV DNA PCR result. (Note: In many cases, there is on-going risk of postnatal transmission through breastfeeding, so an early DNA PCR result indicates infection status at the time of the test, but not the final infection status).
HIV-positive mother	Defined for this survey as mothers whose infants have a positive DBS ELISA.
HIV status unknown	Refers to people (including children) who have not taken an HIV test or who do not know the result of their test.
Infant	A child from birth to 12 months of age.
Infant HIV infection prevalence	Proportion of confirmed HIV-positive (infected) infants among all infants tested during the study period, measured as number of positive DNA PCR

	<p>infant DBS divided by the total number of ELISA samples tested. In this study infant HIV infection prevalence at 6 weeks will be measured in infants age 4 to 8 weeks, who are attending routine immunisation clinic. It will be measured as a point prevalence with the numerator defined as those infants with a positive HIV DNA PCR test and the denominator of all infants tested using HIV ELISA on dried blood spots in the study.</p>
Maternal HIV prevalence	Number of positive (infant) DBS ELISA divided by total number of ELISA samples tested.
Maternal HIV Incidence/Maternal potential HIV acquisition during pregnancy	The number of positive infant DBS ELISA among mothers reporting an HIV-negative status during the interview divided by total number of mothers reporting an HIV-negative status during the interview. This indicator is likely a combination of the following scenarios: (i) mothers who do not wish to admit positive status and report being HIV negative; (ii) mothers who were tested during the window period; (iii) poor quality control or performance of rapid tests in the field causing false negative results at antenatal care (ANC) on HIV-infected women. Reported field sensitivities are as low as 87% to 95% depending on the rapid test used; and (iv) true acquisition of HIV after the last HIV test.
Mother-to-child transmission (MTCT)	Transmission of HIV from an HIV-positive woman to her infant during pregnancy, delivery or breastfeeding. The term is used because the immediate source of the infection is the mother, and does not imply blame on the mother.
MTCT rate	Defined for this survey as a numerator of HIV-positive infants (PCR positive) and denominator of HIV-exposed infants (infant ELISA antibody positive).
Routine offer of counselling and testing	HIV testing that is routinely offered to all ANC clients. Health care personnel provide group information first, followed by individually offering HIV tests. The patient/client has the option to decline testing at any stage of this process. The patient/client receives post-refusal counselling or post-test counselling as appropriate.
Transmission in PMTCT programme	Number of positive DNA PCR and positive ELISA divided by the number of ELISA positive mothers who recall taking ARV prophylaxis or ART during pregnancy/delivery.
Transmission in those not participating in PMTCT Programme (missed opportunities)	Number of positive DNA PCR and positive ELISA divided by number of mothers who do not recall taking ARV prophylaxis or ART during pregnancy/delivery.

## 1. INTRODUCTION

---

The South African National Strategic Plan on HIV and AIDS and STIs (NSP), 2011-2016 aims to reduce HIV transmission from mother to child to less than 2% at six weeks after birth and to less than 5% at 18 months post-delivery by 2016.<sup>4</sup> The NSP acknowledges that strengthening the management, leadership and coordination of the PMTCT programme and ensuring its integration with maternal and child health services is critical.

In 2001 South Africa started implementing a programme to prevent HIV transmission from mother-to-child at 18-pilot sites. The first interventions included single-dose nevirapine (Sd NVP) during labour for the mother and to the baby within 72 hours of delivery; modified obstetric practices; infant feeding counselling and the provision of free commercial infant formula to HIV-positive mothers who avoided breastfeeding (NDOH, 2001).<sup>5</sup> PMTCT interventions were scaled up in 2002 and in 2008 the national antiretroviral regimens for pregnant women were improved to dual therapy (AZT from 28 weeks with Sd-NVP at the outset of labour for pregnant women and Sd-NVP with AZT for baby).<sup>1</sup>

In 2010, PMTCT interventions were further modified as shown in Table 1 (NDOH/SANAC, 2010).<sup>2</sup> The 2010 modifications included routine HIV testing and counselling for pregnant women, dual therapy to prevent MTCT of HIV, ART for pregnant women with CD4 cell count  $\leq 350$  cells/ $\mu$ l, postnatal infant prophylaxis for breastfeeding HIV-positive women and intensified efforts to integrate PMTCT services into routine maternal and child health (MCH) services. These efforts are to meet the NSP targets of reducing the MTCT rate of HIV to less than 5% by 2011 and to meet the 4<sup>th</sup> and 6<sup>th</sup> Millennium Development Goals (MDGs) (i.e., 'reduce by two thirds, between 1990 and 2015, the under-five mortality rate' and 'have halted by 2015 and begun to reverse the spread of HIV/AIDS') (UN, 2011).

The 2011 SAPMTCTE is the first evaluation of PMTCT Option A and was started 16 months after the adoption of the 2010 PMTCT guidelines (PMTCT Option A) and during the month when South Africa changed its infant feeding policy to exclusive breastfeeding for six months and continued breastfeeding thereafter regardless of HIV status following the Breastfeeding Summit.<sup>6</sup> However the provision of free commercial infant formula milk was phased out between August 2011 and April 2012; thus during the 2011 survey free commercial infant formula milk was still provided as part of the PMTCT programme.

The 2011 SAPMTCTE aimed to monitor the 2010-2011 trend in early effectiveness of the South African National PMTCT programme. The primary objective of the 2011 SAPMTCTE was to measure rates of early MTCT of HIV at 4-8 weeks postpartum. The secondary objective was to periodically estimate uptake of key PMTCT interventions and services (e.g., HIV testing, CD4 cell count testing, infant ARV prophylaxis, infant feeding counselling).


## 2. METHODOLOGY

### 2.1 Survey Design and Justification

A cross-sectional facility-based survey, using a biomedical marker to determine MTCT rate, was conducted. The survey was conducted among caregiver-infant pairs who presented at their local primary health care facility for their infant's six-week immunisation (1<sup>st</sup> DTP dose) visit. South Africa reports >95% coverage of six week immunisation (1<sup>st</sup> DTP dose) (WHO, 2011), making these clinics an ideal catchment point for young infants -- of known or unknown HIV exposure status. This provided a convenient sample to determine overall PMTCT effectiveness with relatively limited selection bias.

This methodology has been proven effective in a South African context. Based on the approach recommended by Rollins et al. <sup>7,8</sup> we used a biomedical marker to identify infants exposed to HIV. Chantry et al. <sup>9</sup> found that sero-reversion for ELISA in HIV-exposed infants was not seen prior to 17 weeks of age suggesting that most, if not all, infants aged 4-8 weeks will still have maternal antibodies in their bloodstream. In addition, rapid tests performed on whole blood are less sensitive than HIV ELISA tests so that those performed on similar sample as this study, i.e. small volume of whole blood, would rarely demonstrate sero-reversion before 2 months of age. <sup>10</sup> Therefore, DBS HIV ELISA would be expected to detect the vast majority of HIV-exposed infants. This means that screening infants for the presence of HIV antibody would be a direct measure of infant HIV exposure and an indirect measure of maternal HIV infection prevalence.

Figure 1 Using ELISA as a biomedical marker to identify HIV-exposed infants using HIV antibodies





This evaluation thus aimed to provide:

- 1) A valid estimate of MTCT and HIV infection prevalence in children aged 4-8 weeks, and
- 2) A reasonable estimate of coverage of key PMTCT programme indicators through 6 weeks postpartum.

## **2.2 Study Population and Inclusion/Exclusion Criteria**

The study population comprised infants aged 4-8 weeks and their caregivers visiting public health facilities for the infant's 1<sup>st</sup> DTP dose during the study period.

### ***Inclusion Criteria***

Study participants included 4-8 week old infants attending clinic for 1st DTP immunisation. Caregivers had to consent to participation (consent for maternal or caregiver interview and/or infant DBS).

### ***Exclusion Criteria***

Severely ill infants needing emergency medical care or urgent referral to the next level of care (e.g., infants who are vomiting everything or have convulsions; are lethargic or unconscious; or have severe pneumonia or severe dehydration) were excluded from the study.

## **2.3 Sampling**

### ***Sampling Frame***

The public health facilities were stratified as: < 130, 130-300 and >300 immunisations per year, and data were extracted from the 2007 South African DHIS (Personal Communication C Hedberg, 2009). A strategic decision was made to exclude the small facilities (<130 immunisations per year) from the formal sampling frame. The 2008 national antenatal maternal HIV prevalence estimate of 29% (NDOH, 2009) was used as the cut-off point for classifying facilities as above or below national average for antenatal HIV prevalence.<sup>11</sup> This stratification was only applied to facilities in the large stratum (>300 immunisations per year). A total of 23 strata across province, facility size and maternal HIV prevalence were utilised in the survey sampling frame and were sorted by province, size and maternal HIV prevalence.

### ***Sample Size***

ANC maternal HIV prevalence<sup>11</sup> and estimated MTCT rates from a KwaZulu-Natal survey using similar methodology (Personal communication N. Rollins, unpublished data, 2009) were used to determine the sample size for each province. Specifying relative precisions of 30% to 50% for the expected MTCT rate across provinces plus a design effect of 2 indicated that a total sample size of 12 200 infant DBS specimens were needed. The sample size across provinces ranged from 1 800 (Gauteng) to 700 (Northern Cape).

## **Sampling**

Stratified two-stage sampling was used. In the first stage, facilities (Primary sampling units - PSUs) were randomly sampled proportional to size (PPS) within each stratum. The method operated under the without-replacement-type selection<sup>12</sup> (Lehtonen & Pakhinen, 1994). At the second stage a fixed number of infants per a facility was sampled. The fixed number was the median number of infants expected within the sampling window (three weeks) across the population of facilities within the stratum as determined from the detailed information of the sampling frame above. The fixed number of infants sampled in each facility within a stratum ensured a self-weighting sample. A sampling window of 3 weeks was used to realize the required sample. (Appendix#1)

## **2.4 Data Collection Tools**

Data were gathered using a questionnaire adapted from several validated tools (Rollins et.al., 2007 & 2009; HSRC, undated; Nyblade & MacQuarrie, 2006; Tlebere et.al., 2007; Jackson et al. 2007). The questionnaire included information on maternal age, parity, socio-economic status, antenatal care, HIV testing, maternal HIV status, PMTCT care during pregnancy and delivery, infant feeding counselling, birth information, infant feeding practices, infant weight; immunisations, postnatal visits and illness since birth. Fathers/legal guardians/non-maternal caregivers were administered a shorter form of the questionnaire that excluded ANC and PMTCT information.

The study tool was piloted in the Western Cape and KwaZulu-Natal provinces to test it in English and at least one other official/local languages. Approximately 5-10 participants were administered the study tool in each language as part of the pilot. The primary objective was to test the flow of questions and basic understanding by the participants. The cell phone technology used for data collection, including skips and field data entry, was also examined and tested. Adjustments to the tool and/or cell phone data entry platform were made after the pilot as necessary.

## **2.5 Ethical Considerations**

Written, signed, informed consent for all procedures in the study was obtained from each eligible caregiver for the interview and DBS sampling (separately). Informed consent was in the preferred language of the participants. The information sheet was written in plain lay words that could be easily understood by participants. A confidential Study ID was given to each participant and inserted in consent forms, lab forms and questionnaire for the purpose of data linking and auditing, and to provide the infants' blood test results to mothers or legal guardians. Care was taken to ensure that HIV-infected mothers who refused the study understood that their infant could be tested without participating in the study.

Ethical approval was obtained from the Medical Research Council and from each of the nine provincial research ethics committees. Ethical approval was also granted from the United States Centers for Disease Control and Prevention Atlanta.

## 2.6 Data Collection Methods

Data collection commenced at different times in each province (Table 1). All data collection was completed by 16 March 2012.

**Table 1 Data collection start and end dates in each province**

PROVINCE	2011 Survey		Mop-up - 2011	
	Data collection start date	Data collection end date	Mop-up start date (2012)	Mop-up end date (2012)
<b>Eastern Cape</b>	22 Aug	15 Dec	-	-
<b>Free State</b>	15 Aug	15 Dec	-	-
<b>Gauteng</b>	15 Aug	15 Dec	30 Jan	24 Feb
<b>KwaZulu Natal</b>	15 Aug	15 Dec	13 Feb	16 March
<b>Limpopo</b>	15 Aug	15 Dec	-	-
<b>Mpumalanga</b>	15 Aug	15 Dec	30 Jan	10 March
<b>Northern Cape</b>	15 Aug	15 Dec	-	-
<b>Northwest</b>	15 Aug	15 Dec	-	-
<b>Western Cape</b>	15 Aug	15 Dec	-	-
<b>TOTAL</b>	<b>15 Aug</b>	<b>15 Dec</b>	<b>30 Jan</b>	<b>16 March</b>

### ***Enrolment***

Data collectors recruited mothers/caregivers from the PHC/CHC waiting room during immunisation days. Data collectors introduced themselves and the study verbally and in written form using a standardised information sheet. If the mother agreed to be interviewed, the interview was conducted in a private location. Mother/Infant pairs attending the sampled facilities to receive the infants' DPT first dose vaccination were approached to enroll in the study. A screening questionnaire was administered to determine eligibility and full informed consent forms were completed.

### ***Cell Phone Technology for Data Collection***

Electronic questionnaires were loaded on low-cost mobile phones using the Mobile Researcher software management solution. The Mobile Researcher system consists of three components: the handset, the web interface (data transport system) and web-based research console (Figure 2). The handset is the device on which the questionnaires are entered. Minimum handset functionality is ensured since phone is WAP (Wireless Application Protocol) enabled. The data is transferred via the GPRS (General Packet Radio Services) network using the WAP platform on the mobile phone. The web-based management console is a secure data capture centre that has controlled access.

Questionnaires were uploaded as they were completed to the central web management console and then removed from the phone, while fieldworkers were in an area of mobile reception. In areas where there was no mobile network reception, the questionnaire was stored on the phone until reaching an area with adequate mobile network coverage when data would be automatically uploaded. The questionnaire responses were available on the web-based console every minute, allowing for real-time monitoring of data collection progress and analysis (Figure 3).

Figure 2 Design phase and data collection flow diagram for the cell-phone data collection system





Figure 3 Example of SAPMTCTE Mobile Researcher web-based interface



## 2.7 Laboratory Methods

The National Institute for Communicable Diseases, a division of the South African National Health Laboratory Service (NHLS), conducted the testing. Questionnaires and DBS specimens were linked using unique study identification numbers and lab order numbers. DBS specimens collected from enrolled and consented infants were tested for HIV by means of a laboratory HIV ELISA test (Genscreen HIV antibody assay). In cases where this was reactive (i.e., identified an infant born to an HIV-positive mother), a qualitative HIV PCR (Cobas AmpliPrep/Cobas TaqMan HIV-1 Qual test version 1.0, Roche Diagnostics, Branchburg, NJ) was performed to determine whether the infant was currently HIV-positive. In the case of a known HIV-positive mother, the study DBS specimens and testing replaced the expected routine EID testing. All results were sent to clinic of origin and returned to mother at either 10 or 14 week immunisation visit.

All aspects of the project were carried out according to strict standard operating procedures (SOPs), and testing was conducted under conditions of good laboratory practice. Specimens received in the laboratory were reviewed against the tracking lists/request form for correctness and adequacy of specimens. Each specimen received unique bar-coded identifiers for tracking and data extraction. Rejected specimens were accompanied by a rejection form with specified reasons and referred to field staff for correction. A tracking list of rejected specimens was held by the lab in electronic format. Specimens were tested and results entered into a LIMS (DISA) system; all results had three levels of review.

The algorithm for testing was decided based on the outcome of initial dual ELISA testing. All reactive specimens and every 10<sup>th</sup> non-reactive specimen were tested using a second ELISA, Vironostika (bioMérieux, France). A total of 690 specimens were included in the analysis. The agreement between the two tests was 99.4% and the sensitivity and specificity of the Genscreen assay was 99.7% and 99.2% respectively. Based on these results it was decided that a single ELISA test, Genscreen, be used. All reactive ELISA tests were referred for DNA testing. In the case of a laboratory ELISA equivocal result, HIV PCR testing was performed as a routine.

The procedure for qualitative PCR testing was by automated Ampliprep/Taqman v1.0 technology (Roche). Evaluation of HIV PCR performance on DBS has demonstrated a sensitivity and specificity of 99.7% and 100% respectively.<sup>13</sup> The data extraction of ALL ELISA reactive results was by location code and the referral of spreadsheet to the PCR testing lab. ALL HIV PCR results were extracted and individual reports generated by name of infant for return to the facility where the infant was tested. The reports forms were standardised and had all the required information based on the original request form. All assays used for surveillance were validated and/or verified prior to use, accredited and the performance monitored by proficiency testing. In the case of discordant results between the mother's self-reported result and the laboratory result an algorithm using the two ELISAs, Western blot and PCR was performed on the DBS to exclude lab error or false positive laboratory results.

The data was extracted to exclude personal patient identifiers and emailed to the researchers. The extracted data was in Excel format. Databases were validated and confirmed at two levels before release. The Excel spreadsheet was then merged with the questionnaire database fortnightly. Laboratory data were sent electronically from the laboratory. Tracking logs (study IDs) were used to link questionnaire data and blood test results. The tracking log was managed by the logistics manager.

Prior to the six-week survey, a study was conducted to validate the use of screening and confirmatory third generation ELISAs on DBS. This work was headed by Professors Gayle Sherman and Adrian Puren, and the samples and funding used for this validation study are part of a separate protocol.

## **2.8 Quality Control of Field Work**

Every attempt was made to minimise errors which may result in variation in the collected data contributing to bias in the results. Quality control (QC) was defined as the operational procedures undertaken within the survey, as prescribed by the survey SOPs, to verify that the survey activities (e.g., interviews, obtaining informed consent forms, pre-test counselling, DBS collection, recording data, reporting data) were conducted in accordance with the defined quality standards. The SOPs focused on QC activities done by the field worker, field worker supervisor, quality control officer and the central team. QC activities aimed to improve the quality and validity of the collected data by:

- Identifying factors that may affect the accuracy and reliability of the data and addressing the identified factors;
- Preventing and correcting errors in the collection of data; and
- Ensuring that field activities align with the study SOPs.

## **2.9 Data Management**

Data captured on the phones were protected with a write-only security model. Fieldworkers could modify and review data while the interview was in progress. Captured data was encoded and stored on the device in the Record Management system which ensured that only the Mobile Researcher application could access the data.

The data was transferred securely to the web console, which uses 128-bit strength encryption. Data storage and back up protocols are compliant to enterprise standards and database servers run RAID to ensure redundancy in case of disk failure.

The uploaded data was reviewed daily to ensure that all fieldworkers were submitting responses in accordance to scheduled work plans. The work plans were developed to achieve the required number of DBS per facility and key questions were identified in the database to estimate and track the collection of blood sample progress.

Questionnaire data was maintained by Mobile Researcher and exported to Excel for data analysis. Anonymised laboratory data (Study ID only) were exported to Excel for merging with questionnaire data. Consent verification from hard copy consent forms were entered into Excel and double checked. Interim data analysis was completed during the course of the study. Data from questionnaire, laboratory results and consent verification were all merged and cross-checked. Data without consent verification was not included for analysis. Duplicates and other inconsistencies across data sets were checked and cleaned according to data standards. Out-of-range and data consistency checks were completed as a component of initial data analysis.

## **2.10 Data Analysis**

### ***Sample Realisation***

A total 572 of the 585 sampled clinics were included in final sample. Reasons for non-inclusion included clinic closure (temporary or permanent) or no longer administering immunisations. The overall realisation was 81% with three provinces having low realisation (Northern Cape, Eastern Cape and Limpopo).

### ***Sample Weights***

Sample weights were calculated for the survey to adjust for differential sampling design across provinces and the sample realisation (as outlined above). The data from provinces were weighted by using the proportional distribution of number of life births observed in 2008 for South Africa over provinces. The realisation weights were done at the district or provincial level depending on the sampled size and realisation within strata. For Northern Cape and Eastern Cape the realisation weighting was done at the provincial level. The realisation weights pertain to the per protocol sample size.

A survey analysis was done which took into account the stratification, the different sampling stages and the finite number of PSUs involved. A weighted analysis was done to obtain national estimates as well as provincial estimates. The infant HIV infection prevalence was estimated at the national population level and in the HIV exposed sub-population. These estimates all have 95% confidence intervals. Design effects are also reported. The survey specification and analysis was done in SAS version 9.2. Descriptive statistics of the demographic profile of the participants was done by province and country-wide, accounting for the survey design and realisation.



### 3. RESULTS

#### 3.1 Sample Realisation and Survey Profile

Table 2 indicates the desired and actual sample size for questionnaire plus DBS sample and realised sample size per province and nationally. All but three provinces realised at least 80% of sample.

**Table 2 2010 SAPMTCTE desired and actual sample size by province**

Province	Desired Sample Size (SS)	Actual SS 2010 # (% Desired SS)	Actual SS 2011 #(% Desired SS)
Eastern Cape	1400	776 (55%)	1194 (85%)
Free State	1300	1143 (88%)	1056 (81%)
Gauteng	1800	1735 (96%)	1607 (89%)
KwaZulu-Natal	1400	1224 (87%)	1052 (75%)
Limpopo	1400	1022 (73%)	1070 (76%)
Mpumalanga	1600	1286 (80%)	1210 (76%)
Northern Cape	700	444 (63%)	506 (72%)
North West	1200	1171 (98%)	1037 (86%)
Western Cape	1400	1381 (99%)	1374 (98%)
South Africa	12 200	10 182 (83%)	10106 (83%)

Weighting during analysis adjusted estimates in all provinces with lower than expected sample realisation.

Figure 4 details the final study profile for the survey. Of the 11377 approached at the selected sites 614 (5.40%) did not meet the inclusion criteria, 289 (2.54%) refused participation and 158 (1.39%) had incomplete consent forms. Thus 10475 (92.07%) were enrolled into the SAPMTCT Evaluation. Of these 330 (3.15%) refused infant DBS and 39 (0.37%) had insufficient DBS. Thus 10106 (88.83% of infants screened) had valid interviews and DBS and were included in this analysis




Figure 4 2011 SAPMTCTE study profile

### 3.2 Sample Description and Characteristics

Table 3 provides a summary of selected characteristics of the SAPMTCTE survey sample.

**Table 3 Selected socio-demographic observations of 2011 SAPMTCTE**

**Key:** ZA: South Africa EC: Eastern Cape FS: Free State GP: Gauteng KZN: KwaZulu-Natal  
LP: Limpopo MP: Mpumalanga NC: Northern Cape NW: North West WC: Western Cape

Characteristics	Categories	ZA weighted freq (Wt%) (95% CI)	EC Wt% (95% CI)	FS Wt% (95% CI)	GP Wt% (95% CI)	KZN Wt% (95% CI)	LP Wt% (95% CI)	MP Wt% (95% CI)	NC Wt% (95% CI)	NW Wt% (95% CI)	WC Wt% (95% CI)
Relationship of interview to child	Mother	1156201 (97.02) (96.67-97.36)	94.05 (92.53-95.58)	96.50 (95.66-97.35)	98.44 (97.97-98.92)	96.52 (95.38-97.66)	96.54 (95.54-97.54)	97.65 (96.97-98.34)	98.62 (97.91-99.32)	97.35 (96.64-98.07)	97.29 (96.58-98.01)
	Father	1671 (0.14) (0.07-0.21)	0.08 (0.0-0.23)	0.08 (0.0-0.19)	0.19 (0.0-0.37)	0.07 (0.0-0.18)	0.09 (0.0-0.26)	0.29 (0.02-0.57)	Nil	0.17 (0.0-0.36)	0.21 (0.01-0.40)
	Grandmother/ grandfather	19494 (1.64) (0.37-0.66)	3.85 (2.67-5.03)	2.09 (1.41-2.77)	0.93 (0.52-1.34)	1.66 (0.86-2.45)	1.96 (1.14-2.78)	0.96 (0.52-1.39)	0.59 (0.18-1.01)	1.76 (1.16-2.36)	1.01 (0.53-1.49)
	Guardian/lega l guardian	6171 (0.52) (0.57-0.66)	1.34 (0.72-1.96)	1.16 (0.69-1.63)	0.25 (0.0-0.51)	0.39 (0.082)	0.56 (0.18-0.94)	0.53 (0.16-0.90)	0.79 (0.32-1.26)	0.28 (0.05-0.52)	0.29 (0.06-0.51)
	Other caregiver	8209 (0.67) (0.51-0.87)	0.67 (0.29-1.06)	0.17 (0.0-0.35)	0.19 (0.0-0.37)	1.37 (0.63-2.10)	0.84 (0.34-1.35)	0.57 (0.24-0.91)	Nil	0.44 (0.14-0.74)	1.21 (0.67-1.75)
Age of mother	<15	14800 (1.24) (1.04-1.45)	3.27 (2.41-4.12)	1.23 (0.70-1.77)	0.44 (0.14-0.73)	1.23 (0.58-1.87)	1.02 (0.48-1.58)	1.78 (1.15-2.40)	0.59 (0.17-1.01)	0.66 (0.30-1.01)	1.54 (0.96-2.12)
	15 - 19	185919 (15.6) (14.82-16.38)	21.61 (19.55-23.67)	13.90 (12.08-15.72)	10.89 (9.42-12.36)	19.51 (16.96-22.06)	13.83 (11.81-15.86)	19.92 (17.83-22.00)	15.8 (13.82-17.80)	14.35 (12.34-16.36)	13.47 (11.86-15.08)
	20 - 24	359179 (30.14) (29.16-31.12)	30.90 (28.50-33.30)	31.65 (29.60-33.70)	29.25 (27.28-31.22)	31.3 (28.12-34.50)	31.78 (29.08-34.47)	30.37 (27.70-33.04)	29.05 (26.45-31.65)	31.35 (29.21-33.49)	25.51 (23.50-27.52)

Characteristics	Categories	ZA weighted freq (Wt%) (95% CI)	EC Wt% (95% CI)	FS Wt% (95% CI)	GP Wt% (95% CI)	KZN Wt% (95% CI)	LP Wt% (95% CI)	MP Wt% (95% CI)	NC Wt% (95% CI)	NW Wt% (95% CI)	WC Wt% (95% CI)
Age of mother cont.	25 -29	304662 (25.56) 24.67-26.46)	21.61 19.89-23.32)	25.66 (23.55-27.76)	27.38 (25.53-29.23)	24.79 (21.92-27.65)	25.05 (22.52-27.57)	23.17 (21.12-25.19)	25.89 (23.49-28.28)	25.99 (23.56-28.42)	29.10 (27.01-31.12)
	30 -34	181495 (15.23) (14.63-15.93)	12.48 (11.07-13.89)	17.14 (15.22-19.07)	17.30 (15.76-18.84)	13.31 (11.22-15.41)	15.51 (13.45-17.57)	13.91 (12.28-15.53)	17.59 (15.43-19.74)	13.01 (11.46-14.56)	17.77 (16.08-19.46)
	35 -39	112037 (9.40) (8.80-10.00)	7.70 (6.38-9.03)	8.07 (6.75-9.40)	11.45 (10.04-12.86)	7.32 (5.73-8.90)	9.91 (7.87-11.94)	8.07 (6.79-9.35)	9.29 (8.01-10.57)	11.09 (9.48-12.69)	10.14 (8.94-11.35)
	40 -44	31252 (2.62) (2.31-2.93)	1.76 (0.21-2.25)	2.13 (1.50-2.75)	3.11 (2.42-3.81)	2.24 (1.32-3.17)	2.90 (1.95-3.84)	2.79 (2.03-3.56)	1.78 (1.17-2.39)	3.56 (2.73-4.38)	2.37 (1.69-3.05)
	45 -49	2404 (0.20) (0.11-0.29)	0.67 (0.34-0.10)	0.21 (0.00-0.42)	0.19 (0.01-0.37)	0.29 (0.00-0.59)	nil	nil	nil	nil	0.09 (0.00-0.24)
Education of mother	None	17416 (1.46) (1.23-1.69)	1.68 (1.09-2.26)	0.63 (0.21-1.05)	1.25 (0.81-1.68)	1.27 (0.57-1.97)	1.03 (0.55-1.51)	2.49 (1.58-3.39)	1.98 (1.21-2.75)	3.46 (2.45-4.47)	0.69 (0.24-1.15)
	Grade 1-7	158083 (13.27) (12.36-14.17)	19.93 (16.99-22.88)	11.75 (9.98-13.52)	8.09 (6.61-9.57)	13.77 (10.84-16.69)	11.59 (9.45-13.73)	18.64 (15.67-21.62)	14.03 (11.70-16.36)	21.05 (18.50-23.59)	11.00 (8.89-13.12)
	Grade 8-12	943980 (79.21) (78.02-80.40)	72.78 (69.82-75.73)	85.05 (82.92-87.17)	81.58 (78.69-84.47)	80.93 (77.49-84.37)	79.07 (76.85-81.28)	76.53 (73.46-79.60)	82.02 (79.70-84.33)	70.86 (67.82-73.90)	81.55 (78.98-84.12)

Characteristics	Categories	ZA weighted freq (Wt%) (95% CI)	EC Wt% (95% CI)	FS Wt% (95% CI)	GP Wt% (95% CI)	KZN Wt% (95% CI)	LP Wt% (95% CI)	MP Wt% (95% CI)	NC Wt% (95% CI)	NW Wt% (95% CI)	WC Wt% (95% CI)
Education of mother cont.	Completed tertiary /technical/ university	69924 (5.87) (5.15-6.58)	5.19 (3.90-6.49)	2.30 (1.62-2.98)	2.94 (6.80-11.12)	4.04 (2.63-5.45)	8.13 (6.47-9.79)	2.11 (1.43-2.80)	1.38 (0.61-2.16)	4.48 (3.19-5.78)	6.32 (4.42-8.23)
	Unknown	2344 (0.20) (0.12-0.28)	0.42 (0.12-0.72)	0.28 (0.0-0.55)	0.13 (0.0-0.34)	nil	0.19 (0.0-0.42)	0.23 (0.01-0.46)	0.59 (0.20-0.99)	0.15 (0.0-0.32)	0.45 (0.16-0.74)
Marital status of mother	Single	8851176 (74.28) (72.68-75.87)	73.37 (71.06-75.67)	65.01 (61.86-68.17)	69.07 (64.77-73.38)	89.58 (85.74-93.41)	62.71 (58.72-66.70)	82.04 (79.53-84.54)	80.44 (77.48-83.39)	81.06 (78.36-83.75)	62.30 (57.82-66.77)
	Married	234114 (19.65) (18.49-20.79)	25.63 (23.21-28.05)	27.10 (24.20-30.00)	18.79 (15.93-21.66)	7.33 (5.61-9.04)	32.06 (28.24-35.87)	15.45 (12.82-18.08)	17.98 (14.83-21.14)	16.57 (14.13-19.01)	29.23 (25.61-32.85)
	Co-habiting	66914 (5.62) (4.27-6.96)	0.67 (0.26-1.08)	6.98 (5.02-8.94)	11.51 (7.53-15.49)	3.10 (0.0-6.82)	4.95 (2.83-7.08)	2.24 (1.36-3.13)	0.99 (0.12-1.86)	1.75 (1.06-2.44)	7.27 (4.95-9.60)
	widowed	2079 (0.17) (0.11-0.24)	nil	0.63 (0.12-1.15)	0.25 (0.04-0.45)	nil	nil	0.17 (0.0-0.37)	0.59 (0.18-1.00)	0.45 (0.12-0.79)	0.23 (0.02-0.42)
	Divorced / separated	3269 (0.27) (0.18-0.37)	0.34 (0.07-0.60)	0.28 (0.0-0.55)	0.31 (0.02-0.60)	nil	0.28 (0.01-0.55)	0.10 (0.0-0.28)	nil	0.17 (0.0-0.37)	0.98 (0.60-1.37)
	Unknown	195.13 (0.02)	nil	nil	0.06 (0.0-0.17)	nil	nil	nil	nil	nil	nil

Characteristics	Categories	ZA weighted freq (Wt%) (95% CI)	EC Wt% (95% CI)	FS Wt% (95% CI)	GP Wt% (95% CI)	KZN Wt% (95% CI)	LP Wt% (95% CI)	MP Wt% (95% CI)	NC Wt% (95% CI)	NW Wt% (95% CI)	WC Wt% (95% CI)
Main building material of house	Brick/Cement block	902267 (75.74) (73.77—77.71)	59.06 (52.77-65.35)	79.71 (77.12-82.30)	80.52 (76.62-84.43)	68.29 (61.94-74.64)	87.45 (84.79-90.12)	89.90 (87.25-92.54)	79.25 (75.27-83.23)	74.51 (71.25-77.76)	70.26 (64.11-76.42)
	Informal material / corrugated iron / wood	190680 (16.01) (14.52-17.49)	9.90 (6.64-13.16)	18.96 (16.44-21.47)	19.48 (15.57-23.38)	12.38 (9.03-15.74)	7.96 (5.80-10.12)	7.95 (5.54-10.36)	20.75 (16.77-24.73)	23.46 (20.18-26.74)	29.74 (23.59-35.90)
	Traditional material/mud	97799 (8.21) (6.43-9.98)	31.04 (23.55-38.53)	1.33 (0.54-2.13)	No data	19.13 (12.03-26.24)	4.59 (2.48-6.69)	2.09 (0.88-3.30)	nil	2.03 (0.89-3.16)	nil
	Other	529.76 (0.05) (0-0.09)	nil	nil	nil	0.20 (0.0-43)	nil	0.06 (0.0-1.6)	nil	nil	nil
Main source of drinking water	Piped in house or yard	872976 (73.28) (70.84-75.72)	46.56 (38.78-54.35)	96.52 (95.63-97.41)	92.91 (90.53-95.28)	55.34 (46.34-64.33)	42.88 (37.87-47.90)	85.37 (80.49-90.25)	90.32 (87.60-93.04)	74.83 (68.94-80.73)	98.01 (96.90-99.12)
	Not piped in house or yard	318299 (26.72) (24.28-29.16)	53.44 (45.66-61.22)	3.48 (2.59-4.37)	7.09 (4.72-9.47)	44.66 (35.67-53.66)	57.12 (52.10-62.14)	14.63 (9.75-19.51)	9.68 (6.97-12.40)	25.17 (19.27-31.06)	1.99 (0.88-3.10)
Type of toilet	Flush toilet	598233 (50.22) (47.85-52.58)	24.58 (18.28-30.88)	72.00 (66.94-77.05)	87.06 (83.50-90.62)	22.56 (16.11-29.01)	13.76 (9.17-18.36)	20.80 (14.92-26.69)	80.63 (76.72-84.54)	43.58 (35.75-51.40)	92.61 (90.01-95.22)
	Pit latrine	554604 (46.56) (44.24-48.87)	69.13 (62.88-75.38)	25.43 (20.48-30.39)	12.63 (9.10-16.16)	71.04 (64.12-77.95)	81.84 (77.22-86.45)	78.88 (73.00-84.75)	11.86 (9.22-14.50)	54.31 (46.59-62.04)	5.26 (2.83-7.69)

Characteristics	Categories	ZA weighted freq (Wt%) (95% CI)	EC Wt% (95% CI)	FS Wt% (95% CI)	GP Wt% (95% CI)	KZN Wt% (95% CI)	LP Wt% (95% CI)	MP Wt% (95% CI)	NC Wt% (95% CI)	NW Wt% (95% CI)	WC Wt% (95% CI)
Type of toilet Cont.	None	34037 (2.86) (1.77-3.94)	5.87 (3.69-8.05)	0.55 (0.16-0.94)	0.19 (0.01-0.37)	6.41 (1.37-11.44)	4.31 (2.69-5.92)	0.32 (0.07-0.57)	3.36 (2.00-4.72)	2.02 (1.27-2.78)	1.20 (0.57-1.83)
	Other	4401 (0.37) (0.25-0.48)	0.42 (0.05-0.79)	2.02 (0.68-3.36)	0.13 (0.0-0.34)	nil	0.09 (0.0- 0.26)	nil	4.15 (2.49- 5.81)	0.09 (0.0-0.24)	0.93 (0.46- 1.40)
Main source of fuel	Electricity/ gas/ paraffin	1064496 (89.36) (87.47-91.24)	88.84 (83.41-94.27)	98.74 (97.90-99.57)	99.25 (98.70-99.81)	81.48 (74.23-88.72)	57.21 (49.56-64.86)	97.35 (96.05-98.65)	96.25 (94.60-97.89)	96.00 (94.77-97.24)	99.56 (99.28-99.84)
	Other	126779 (10.64) (8.76-12.52)	11.16 (5.73-16.59)	1.26 (0.43-2.10)	0.75 (0.20-1.30)	18.52 (11.28-25.77)	42.79 (35.14-50.44)	2.65 (1.35-3.95)	3.76 (2.11-5.40)	4.00 (2.77-5.23)	0.44 (0.16-0.72)
Depletion of food supply in past 12 months	Yes	153521 (12.89) (11.58-14.20)	12.42 (9.08-15.75)	14.90 (12.77-17.03)	8.15 (5.92-10.39)	18.35 (13.48-23.22)	17.14 (13.97-20.30)	5.43 (3.55-7.31)	21.15 (18.46-23.84)	10.81 (8.54-13.08)	14.80 (12.45-17.45)
	No	1035236 (86.90) (85.59-88.22)	87.16 (83.77-90.56)	84.93 (82.70-87.16)	91.72 (89.48-93.97)	81.55 (76.66-86.44)	82.87 (79.70-86.03)	94.44 (92.52-96.37)	78.66 (75.90-81.41)	88.54 (86.29-90.80)	84.72 (82.04-87.40)
	Don't know	2519 (0.21) (0.13-0.29)	0.42 (0.10-0.74)	0.17 (0.0-0.41)	0.13 (0.0-0.27)	0.10 (0.0-0.28)	nil	0.13 (0.0-0.26)	0.20 (0.0-0.45)	0.65 (0.26-1.03)	0.48 (0.17-0.79)
Planned Pregnancy	Yes	437343 (38.37) (36.83-39.90)	31.99 (27.09-36.88)	50.16 (46.50-53.81)	43.12 (39.57-46.67)	22.67 (19.34-26.01)	52.61 (49.88-55.35)	47.15 (41.63-52.66)	38.71 (34.54-42.88)	34.46 (31.44-37.48)	38.36 (35.76-40.95)
Planned Pregnancy	No	699693 (61.38) (59.84-62.92)	67.74 (62.89-72.59)	48.78 (44.78-52.79)	56.75 (53.19-60.32)	77.02 (73.68-80.36)	47.39 (44.66-50.12)	52.77 (47.26-58.27)	60.89 (56.68-65.10)	64.73 (61.68-67.77)	61.64 (59.05-64.24)

Characteristics	Categories	ZA weighted freq (Wt%) (95% CI)	EC Wt% (95% CI)	FS Wt% (95% CI)	GP Wt% (95% CI)	KZN Wt% (95% CI)	LP Wt% (95% CI)	MP Wt% (95% CI)	NC Wt% (95% CI)	NW Wt% (95% CI)	WC Wt% (95% CI)
cont.	DKN	2849 (0.25) (0.15-0.35)	0.28 (0.01-0.54)	1.06 (0.20-1.92)	0.13 (0.0-0.28)	0.31 (0.0-0.62)	nil	0.09 (0.0-0.22)	0.40 (0.06-0.75)	0.82 (0.27-1.37)	nil
Infant gender	Male	596132 (50.02) (49.04-51.00)	51.84 (49.24-54.45)	50.36 (47.61-53.11)	50.97 (48.89-53.04)	49.66 (46.54-52.78)	48.04 (45.63-50.44)	50.38 (48.54-52.22)	54.15 (51.53-56.77)	46.81 (44.46-49.15)	49.35 (47.33-51.37)
	Female	595615 (49.98) (49.00- 50.95)	48.16 (45.56-50.76)	49.64 (46.89-52.39)	49.04 (46.96-51.11)	50.35 (47.23-53.47)	51.96 (49.56-54.37)	49.62 (47.78-51.46)	45.85 (43.23-48.47)	53.19 (50.85-55.54)	50.65 (48.63-52.67)
Infant age in weeks	4	12158 (1.02) (0.80-1.24)	3.09 (2.10-4.10)	0.54 (0.17-0.91)	0.25 (0.04-0.46)	1.05 (0.31-1.79)	0.65 (0.18-1.13)	3.03 (2.18-3.88)	0.59 (0.20-0.98)	0.56 (0.21-0.92)	0.07 (0.00-0.18)
	5	74632 (6.26, 5.40-7.12)	10.97 (7.91-14.03)	8.72 (6.06-11.39)	3.24 (2.01-4.46)	6.60 (3.68-9.52)	6.54 (4.03-9.05)	9.31 (7.28-11.34)	9.09 (6.73-11.45)	8.07 (6.178-9.96)	2.18 (1.19-3.16)
	6	95558 (80.18, 78.93-81.43)	64.57 (60.62-68.53)	82.60 (79.40-85.79)	86.43 (83.76-89.11)	79.09 (75.61-82.56)	86.63 (83.58-89.69)	71.51 (67.91-75.10)	74.11 (70.54-77.68)	79.53 (77.26-81.81)	83.79 (81.40-86.19)
	7	120393 (10.10, 9.25-10.95)	16.50 (13.76-19.23)	7.57 (5.94-9.20)	8.46 (6.37-10.56)	10.90 (8.74-13.06)	4.67 (3.01-6.34)	12.14 (10.12-14.15)	13.24 (11.50-14.98)	9.44 (7.74-11.15)	11.27 (9.22-13.33)
	8	29006 (2.43) (2.11- 2.75)	4.86 (3.49-6.22)	0.56 (0.28-0.85)	1.62 (1.05-2.19)	2.36 (1.50-3.23)	1.50 (0.76-2.23)	4.02 (2.80-5.23)	2.96 (1.92-3.29)	2.38 (1.47-3.29)	2.68 (1.85-3.51)



Of note is that nationally 97.02% of infants were brought to the clinic by their mothers; 91.93% mothers were aged 20-34 years; 85.08% mothers had completed grades 8-12 or more of school; 74.28% of mothers were single, 61.38% of mothers reported that their pregnancy was unplanned, 12.89% reported running out of food at some time during the past 12 months and 91.7% of infants were aged 4-8 weeks

### 3.3 Infant HIV Infection Prevalence

**Table 4 Weighted Infant HIV infection prevalence nationally and by province**

Province	% Infant HIV Infection Prevalence 2010	% Infant HIV Infection Prevalence 2011
<b>Eastern Cape</b>	2.0 (1.1-2.9)	1.3 (0.7-1.8)
<b>Free State</b>	2.4 (1.6-3.2)	1.2 (0.7-1.7)
<b>Gauteng</b>	1.1 (0.6-1.5)	0.8 (0.3-1.2)
<b>KwaZulu-Natal</b>	1.9 (1.2-2.7)	0.9 (0.4-1.5)
<b>Limpopo</b>	0.9 (0.4-1.5)	0.8 (0.3-1.2)
<b>Mpumalanga</b>	3.0 (2.1-3.8)	1.2 (0.8-1.7)
<b>Northern Cape</b>	0.3 (0.1-0.6)	1.0 (0.4-1.6)
<b>Northwest</b>	1.9 (1.2-2.5)	0.8 (0.4-1.2)
<b>Western Cape</b>	0.9 (0.4-1.5)	0.4 (0.1-0.6)
<b>National</b>	1.5 (1.3-1.7)	0.9 (0.7-1.1)

The national weighted infant HIV infection prevalence among infants aged 4-8 weeks attending child health clinics for their six week immunisation was 0.9% (95%CI: 0.7-1.1%). (Table 4) Infant HIV infection prevalence is the rate of HIV-positivity among all infants tested regardless of exposure which provides an indication of total burden of HIV disease in infants at 4-8 weeks of age.

### 3.4 National and Provincial Infant HIV Exposure and MTCT Rates

The national rate of infant HIV exposure was 32.2% (95%CI: 30.7-33.6%), with wide provincial variation. (Table 5) (Note: Infant HIV exposure prevalence is presumed to be roughly equivalent to maternal HIV prevalence.)

Among these HIV-exposed infants, the national rate of MTCT of HIV by 8 weeks is 2.7% (95%CI: 2.1-3.2%), with an almost 3-fold difference between provinces; the lowest rate of 1.98% (95%CI: 0.65-3.31) was found in the Western Cape and the highest rate of 6.1% (95%CI: 2.5-9.6) in the Northern Cape.

It is important to note that for the Eastern Cape and Northern Cape (\*) provinces in 2010 and the Northern Cape province in 2011, the point estimates are correct but the sample precision was less (wider confidence intervals). This was due to the lower sample realisation rates,

**Table 5 Weighted infant HIV exposure and 4-8 week (early) MTCT of HIV by province for 2010 and 2011**

Province	Infant HIV exposure 2010 (%)	MTCT (%) 95% CI 2010	Infant HIV exposure 2011 (%)	MTCT (%) 95% CI 2011
Eastern Cape*	30.5 (26.9-34.2)	4.7 (2.4-7.0)	32.0 (29.6-35.5)	3.82 (2.1-5.5)
Free State	31.3 (29.1-33.5)	5.9 (3.8-8.0)	30.9 (28.6-33.3)	3.80 (2.3-5.3)
Gauteng	30.4 (27.9-33.0)	2.5 (1.5-3.6)	33.1 (29.8-36.4)	2.13 (0.2-3.4)
KwaZulu-Natal	44.3 (40.2-48.4)	2.9 (1.7-4.0)	44.4 (39.8-48.9)	2.10 (0.9-3.3)
Limpopo	23.9 (21.8-25.9)	3.6 (1.4-5.8)	23.0 (19.9-26.2)	3.06 (1.2-4.9)
Mpumalanga	37.0 (34.3-39.7)	5.7 (4.1-7.3)	35.6 (33.3-37.8)	3.32 (2.2-4.5)
Northern Cape*	16.0 (13.7-18.3)	1.4 (0.1-3.4)	15.1(12.7-17.5)	6.1 (2.5-9.6)
Northwest	31.3 (29.0-33.5)	4.4 (2.9-5.9)	30.8 (28.5-33.1)	2.6 (1.1-4.0)
Western Cape	21.0 (17.0-25.0)	3.9 (1.9-5.8)	17.8 (14.8-20.8)	1.98 (0.65-3.31)
<b>South Africa</b>	<b>32.0 (30.7-33.3)</b>	<b>3.5 (2.9-4.1)</b>	<b>32.2 (30.7-33.6)</b>	<b>2.7 (2.1-3.2)</b>

### 3.5 National PMTCT Programme Cascade

Table 6 presents results for PMTCT programme indicators as per maternal report in all mothers interviewed. The percent of pregnant women with unknown HIV status prior to their first antenatal booking who had an HIV test during pregnancy was 98.3. Maternal receipt of HIV test results was also high at 99.4%. Of ALL mothers enrolled in the survey 29.6% reported being HIV-positive while HIV antibody was found in 32.2% of ALL infants – a 2.9% difference. Of concern is that of those *mothers who reported being HIV-negative*, 3.9% of their infants had HIV antibodies, suggesting a high rate of maternal potential acquisition of HIV infection during pregnancy. This rate also varied substantially across provinces from a low of 0.7% in the Western Cape to a high of 10.2% in Mpumalanga and. The indicator ‘Maternal potential HIV acquisition’ is a likely combination of the following scenarios:

- (i) Mothers did not want to admit being HIV-positive and instead, reported being HIV negative. However, the 2011 data show that refusals for infant HIV testing were low and disclosure was high; thus the contribution that this scenario makes to the indicator is probably minimal.
- (ii) Mothers were tested during the window period for the ANC test.

- (iii) Poor QC/performance of rapid tests in the field causes false negative results at ANC on HIV-infected women. Reported field sensitivities are as low as 87% to 95% depending on the rapid test used. In correlation to this was mothers who reported being HIV-positive but for which infant test HIV antibody negative which was 2.5% (95% CI 1.8-3.2%) and also suggests potential problems with performance of rapid tests during ANC.
- (iv) True acquisition of HIV after the last HIV test – primarily during pregnancy.

**Table 6 HIV testing & results among pregnant women (weighted analysis) for 2010 and 2011**

Province	% ANC HIV Test		% Tested who received result		% Mothers report being HIV-positive		%Infants of reported HIV-negative mothers who had HIV antibody	
	2010	2011	2010	2011	2010	2011	2010	2011
Eastern Cape	97.5 (96.5-98.6)	98.3 (97.6-98.9)	98.1 (97.1-99.1)	98.4 (97.7-99.2)	27.1 (23.5-30.7)	29.9 (26.8-32.9)	7.8 (5.8-9.7)	5.2 (3.7-6.7)
Free State	98.8 (98.3-99.2)	97.6 (96.8-98.5)	98.9 (98.5-99.4)	99.0 (98.5-99.6)	27.9 (25.7-30.1)	28.5 (25.8-31.1)	5.4 (4.3-6.4)	4.2 (3.0-5.3)
Gauteng	99.1 (98.7-99.2)	98.5 (98.0-99.1)	99.3 (98.9-99.6)	99.7 (99.5-99.9)	28.3 (25.8-30.8)	30.4 (27.1-33.7)	3.0 (2.2-3.9)	3.4 (2.4-4.4)
KwaZulu-Natal	98.9 (98.3-99.2)	97.9 (96.8-99.1)	99.5 (99.1-99.9)	99.9 (99.7-100.0)	42.2 (38.1-46.2)	41.4 (36.3-46.4)	3.2 (2.1-4.4)	5.0 (3.7-6.4)
Limpopo	98.6 (97.8-99.5)	98.4 (97.7-99.0)	97.0 (95.9-98.1)	99.4 (98.9-99.8)	19.4 (17.3-21.6)	20.2 (17.3-23.1)	5.1 (3.6-6.7)	1.6 (0.9-2.3)
MP	98.6 (97.8-99.3)	98.4 (97.6-99.1)	97.1 (96.3-98.0)	99.0 (98.6-99.5)	32.6 (29.7-35.5)	29.1 (26.7-31.4)	7.8 (5.8-9.7)	10.2 (8.2-12.2)
Northern Cape	99.3 (98.9-99.8)	99.0 (98.4-99.5)	96.7 (95.7-97.6)	99.8 (99.5-100.0)	14.4 (12.2-16.7)	14.3 (12.3-16.2)	2.2 (1.2-3.3)	1.9 (1.2-2.7)
North West	99.2 (98.8-99.6)	99.1 (98.7-99.6)	98.5 (97.8-99.1)	99.1 (98.5-99.6)	28.7 (26.7-30.6)	29.2 (26.7-31.7)	5.4 (3.9-6.8)	3.7 (2.5-4.9)
Western Cape	98.6 (97.9-99.3)	97.7 (97.1-98.4)	98.8 (98.3-99.3)	99.4 (99.0-99.8)	19.9 (16.1-23.8)	17.2 (14.1-20.3)	1.1 (0.3-1.9)	0.7 (0.3-1.2)
South Africa	98.8 (98.5-99.0)	98.3 (98.0-98.6)	98.6 (98.4-98.9)	99.4 (99.3-99.6)	29.4 (28.1-30.7)	29.6 (28.0-31.2)	4.1 (3.7-4.6)	3.9 (3.5-4.4)


Table 7 shows PMTCT programme indicators for women who reported being HIV-positive: 77.4% of mothers reported getting a CD4 test result; taking ART was reported by 41.9% of mothers, while 52.0% reported receiving both maternal and neonatal antiretroviral (ARV) prophylaxis. This suggests that 93.9% of HIV-positive mothers received either ART or ARV prophylaxis (**Figure 5**). These figures show an increase in ART access compared with 2010 data, probably due to the CD4 cell cut-off for ART access ( $\leq 350$  cells/ $\mu$ l) in 2010. Maternal knowledge of CD4 cell count results remained the same in 2010 and 2011, possibly illustrating the lack of communication between health care personnel and HIV positive women or poor maternal memory of CD4 cell count result.

**Table 7 Access to the PMTCT programme in reported HIV-positive mothers (weighted analysis), 2010 and 2011**

Province	% Received CD4 Test		% Received ARV/ART		% Mother & Infant Received ARV Prophylaxis		% Intended to obtain EID @ 6 weeks	
	2010	2011	2010	2011	2010	2011	2010	2011
EC	67.6 (60.2-75.1)	70.1 (64.1-76.0)	23.0 (16.9-29.0)	38.9 (33.3-44.4)	63.5 (55.3-71.7)	53.5 (48.1-58.9)	21.6 (14.9-28.4)	28.0 (17.6-38.4)
FS	85.8 (82.7-89.0)	63.2 (56.9-69.4)	37.7 (33.2-42.2)	44.2 (38.8-49.7)	56.4 (51.6-61.1)	51.8 (46.5-57.2)	43.7 (33.3-54.1)	24.6 (17.8-31.4)
GP	74.6 (69.8-79.4)	77.1 (72.1-82.0)	40.1 (34.9-45.3)	46.1 (41.1-51.1)	52.8 (47.1-58.4)	48.1 (42.8-53.4)	42.5 (32.6-52.4)	25.4 (19.2-31.5)
KZN	85.5 (82.1-88.8)	87.2 (81.3-93.1)	29.4 (25.5-33.3)	39.0 (34.2-43.8)	65.2 (61.1-69.3)	56.6 (51.8-61.3)	41.1 (30.5-51.6)	63.6 (53.7-73.5)
LP	68.3 (61.0-75.5)	68.1 (62.2-74.0)	33.3 (27.3-39.4)	37.7 (31.9-43.5)	54.3 (47.3-61.3)	48.2 (42.5-53.9)	28.4 (20.4-36.5)	31.1 (23.2-38.9)
MP	69.5 (65.5-73.5)	66.6 (62.6-70.7)	27.5 (23.3-31.7)	31.9 (25.9-37.8)	56.1 (51.8-60.3)	60.5 (54.3-66.7)	29.8 (23.1-36.5)	41.2 (32.1-50.3)
NC	88.7 (83.0-94.3)	76.8 (69.4-84.3)	28.6 (19.6-37.6)	52.2 (46.6-57.7)	58.7 (51.1-66.3)	46.4 (41.3-51.5)	1.6 (0.1-4.0)	11.6 (5.2-18.0)
NW	81.7 (78.3-85.1)	74.2 (69.9-78.5)	33.7 (29.1-38.4)	49.3 (44.6-53.9)	57.4 (52.4-62.5)	44.0 (39.6-48.5)	3.6 (1.6-5.7)	13.0 (7.3-18.7)
WC	89.6 (86.8-92.5)	86.4 (81.7-91.2)	34.2 (27.9-40.6)	49.8 (45.4-54.3)	60.0 (52.7-67.3)	47.6 (43.8-51.4)	37.9 (28.8-47.0)	46.3 (36.1-56.5)
ZA	78.3 (76.4-80.4)	77.4 (74.9-80.0)	33.1 (30.8-35.3)	41.9 (39.7-44.2)	58.7 (56.3-61.1)	52.0 (49.7-54.2)	35.1 (30.6-39.6)	38.5 (34.3-42.7)

Only 38.46% of reported HIV-positive mothers indicated that they planned to obtain early infant diagnosis (EID) for their infant during their six week immunisation visit (ranging from 11.59% in Northern Cape to 63.56% in KwaZulu Natal ).

Figure 5 PMTCT service uptake (PMTCT cascade) in South Africa



**Note:** The first three indicators apply to all mothers, while the last three apply only to those who reported being HIV-positive. Red indicates the percentage receiving ART while blue indicates the percentage receiving ARV prophylaxis.

## 3.6 Demographic Characteristics, MTCT and the PMTCT Cascade by Province

### 3.6.1 Eastern Cape

Eastern Cape achieved a sample realisation of 55%. This was due to a high number of medium size clinics requiring extended resources. Furthermore in the Eastern Cape many infants presenting for six week immunisation were older than 8 weeks of age. This was especially the case in large urban clinics. These infants were not eligible for enrolment into the survey.

#### *General Description of Provincial Sample*

Table 8 presents characteristics of respondents in the Eastern Cape province. Similar to the national trend, the majority of the respondents are single (73.37%) mothers (94.05%), with education level of grade 8-12 (72.78%). In comparison with other provinces, Eastern Cape has a significant percentage (12.42) of respondents that reported experiencing depletion of food in the household in the last 12 months. Economic status indicators also show that pit latrines (69.13%) and not piped water (53.44%) are utilised by the majority of the respondents.

**Table 8 Baseline characteristics of Eastern Cape SAPMTCTE survey participants**

Characteristics	2010		
	Categories	%	95% CI
Relationship to child	Mother	97.3	93.9-96.8
	Caregiver	4.7	3.2-6.1
Mean age of mother (range)	25.1 (14-52)		
Infant gender	Male	52.1	48.7-55.5
	Female	47.9	44.5-51.3
Education of mother	None	2.5	1.6-3.4
	Grade 1-7	21.8	18.4-25.1
	Grade 8-12	70.8	67.4-74.3
	Above Grade 12	4.4	3.0-5.9
Marital status of mother	Single	75.8	73.1-78.6
	Married/cohabitating	23.8	21.1-26.6
Main building material of house	Brick/Cement block	63.1	55.5-70.7
	Informal material	11.8	8.1-15.5
	Traditional material/mud	25.1	18.6-31.5
Main source of drinking water	Piped in house or yard	42.3	34.8-49.8
	Not piped in house or yard	57.7	50.2-65.2
Type of toilet	Flush toilet	26.4	19.6-33.3
	Pit latrine	62.9	56.6-69.1
	None	9.6	5.9-13.4
	Other	1.0	0.4-1.7
Main source of fuel	Electricity/gas/paraffin	97.8	96.9-98.8
	Other	2.2	1.2-3.1
Depletion of food supply in past 12 months	Yes	24.5	18.9-30.1
	No	75.0	69.4-80.6

2011			
Characteristics	Categories	%	95% CI
Relationship to child	Mother	94.0	92.53-95.58
	Father	0.1	0.0-2.3
	Grandmother/grandfather	3.8	2.7-5.0
	Guardian/legal guardian	1.3	0.7-2.0
	Caregiver	0.7	0.3-1.1
Mean age of mother (range)			
Infant gender	Male	51.8	49.2-54.4
	Female	48.2	45.6-50.8
Education of mother	None	1.7	1.1-2.3
	Grade 1-7	19.9	17.0-22.9
	Grade 8-12	72.8	69.8-75.7
	Completed tertiary/technical /university	5.2	3.9-6.5
	Don't know	0.4	0.1-0.7
Marital status of mother	Single	73.4	71.1-75.7
	Married	25.6	23.2-28.1
	Co-habiting	0.7	0.3-1.1
	Widowed	Nil	
	Divorced/separated	0.3	0.1-0.6
	Don't know	Nil	
Main building material of house	Brick/Cement block	59.1	52.8-65.3
	Informal material/corrugated iron/wood	9.9	6.6-13.2
	Traditional material/mud	31.0	23.6-38.5
	other	Nil	
Main source of drinking water	Piped in house or yard	46.6	38.8-54.3
	Not piped in house or yard	53.4	45.7-61.2
Type of toilet	Flush toilet	24.6	18.3-30.9
	Pit latrine including ventilated pit latrine	69.1	62.9-75.4
	None	5.9	3.7-8.1
	Other	0.4	0.1-0.8
Main source of fuel	Electricity/gas/paraffin	88.8	83.4-94.3
	Other	11.2	5.7-16.6
Depletion of food supply in past 12 months	Yes	12.4	9.1-15.8
	No	87.2	83.8-90.6
	Don't know	0.4	0.1-0.7
Was this pregnancy planned	Yes	32.0	27.1-36.9
	No	67.7	62.9-72.6
	Don't know	0.3	0.0-0.5

### Infant HIV Exposure and MTCT Rate in Eastern Cape Province

Text box 1 shows that infants' HIV exposure was 32.0%, with a 1.3% early infant HIV infection prevalence and a 3.8% (95%CI: 2.1-5.5) MTCT rate at 4-8 weeks. The larger confidence interval attached to this estimate is due to the smaller sample size attained in Eastern Cape. The percent of reported HIV-negative mothers whose infants had HIV antibodies (presumed maternal HIV acquisition after the initial HIV test) was 5.2% (95% CI 3.7-6.7%), was significantly lower than in 2010, but still the second highest in South Africa (after Mpumalanga).


**Text Box 1: Eastern Cape infant HIV exposure and MTCT**

2010			
Infant HIV Exposure % (95%CI)	Infant HIV infection prevalence at 4-8 weeks	MTCT @ 4-8 weeks:%(95%CI)	%Infants of reported HIV-negative mothers who had HIV antibody
30.5 (26.9-34.2)	2.0 (1.1-2.9)	4.7 (2.4-7.0)	7.8 (5.8-9.7)
2011			
Infant HIV Exposure % (95%CI)	Infant HIV infection prevalence at 4-8 weeks	MTCT @ 4-8 weeks:%(95%CI)	%Infants of reported HIV-negative mothers who had HIV antibody
32.0 (29.6-35.5)	1.3 (0.7-1.8)	3.8 (2.1-5.5)	5.2 (3.7-6.7)

### PMTCT Service Uptake (PMTCT Cascade) in the Eastern Cape Province

Figure 6 indicates that Eastern Cape has a fairly high antenatal HIV testing rate (97.5%) but ARV prophylaxis/ART coverage of 86.5%. Coverage of CD4 count and intended EID is low in Eastern Cape.

**Figure 6 PMTCT service uptake (PMTCT cascade) in the Eastern Cape**





**Note:** The first three indicators apply to all mothers, while the last three apply only to those who reported being HIV-positive. Red indicates the percentage receiving ART while blue indicates the percentage receiving ARV prophylaxis.

### 3.6.2 Free State

The survey attained 88% of targeted sample size in the Free State.

#### *General Description of Provincial Sample*

In the Free State 65.01% of mothers were single. More than 95% had piped water and electricity, gas or paraffin fuel source, while only 72% had a flush toilet. (Table 9)

**Table 9** Baseline characteristics of Free State SAPMTCTE survey participants

2010			
Characteristics	Categories	%	95% CI
Relationship to child	Mother	96.9	96.2-97.7
	Caregiver	3.1	3.2-6.1
Mean age of mother – mean (range)	25.8 (14-48)		
Infant gender	Male	51.6	49.6-53.6
	Female	48.4	46.4-50.4
Education of mother	None	0.9	0.5-1.4
	Grade 1-7	15.1	13.3-16.8
	Grade 8-12	79.4	77.5-81.3
	Above Grade 12	3.7	2.7-4.6
Marital status of mother	Single	63.5	61.0-66.0
	Married/cohabitating	36.0	33.5-38.5
Main building material of house	Brick/Cement block	78.6	76.3-80.9
	Informal material	19.8	17.7-22.0
	Traditional material/mud	1.5	0.9-2.1
Main source of drinking water	Piped in house or yard	85.1	81.5-88.8
	Not piped in house or yard	14.9	11.2-18.5
Type of toilet	Flush toilet	66.4	60.9-72.0
	Pit latrine	31.1	25.5-36.6
	None	0.2	0.0-0.3
	Other	2.3	1.3-3.4
Main source of fuel	Electricity/Gas/Paraffin	97.3	96.1-98.6
	Other	2.7	1.4-3.9
Depletion of food supply in past 12 months	Yes	13.7	11.6-15.8
	No	86.1	84.0-88.2

2011			
Characteristics	Categories	%	95% CI
Relationship to child	Mother	96.5	95.7-97.3
	Father	0.1	0.0-0.2
	Grandmother/grandfather	2.1	1.4-2.8
	Guardian/legal guardian	1.2	0.7-1.6
	Caregiver	0.12	0.0-0.35
Mean age of mother (range)			
Infant gender	Male	50.4	47.6-53.1
	Female	49.6	46.9-52.4
Education of mother	None	0.6	0.2-1.0
	Grade 1-7	11.8	10.0-13.5
	Grade 8-12	85.1	82.9-87.2
	Completed tertiary/technical /university	2.3	1.6-3.0
	Don't know	0.3	0.0-0.6
Marital status of mother	Single	65.0	61.9-68.2
	Married	27.1	24.2-30.0
	Co-habiting	7.0	5.0-8.9
	Widowed	0.6	0.1-1.2
	Divorced/separated	0.3	0.0-0.6
	Don't know	Nil	
Main building material of house	Brick/Cement block	79.7	77.1-82.3
	Informal material/corrugated iron/wood	19.0	16.4-21.5
	Traditional material/mud	1.3	0.5-2.1
	other	Nil	
Main source of drinking water	Piped in house or yard	96.5	95.6-97.4
	Not piped in house or yard	3.5	2.6-4.4
Type of toilet	Flush toilet	72.0	66.9-77.0
	Pit latrine including ventilated pit latrine	25.4	20.5-30.4
	None	0.6	0.2-0.9
	Other	2.0	0.7-3.4
Main source of fuel	Electricity/gas/paraffin	98.7	97.9-99.6
	Other	1.3	0.4-2.1
Depletion of food supply in past 12 months	Yes	14.9	12.8-17.0
	No	84.9	82.1-87.2
	Don't know	0.2	0.0-0.4
Was this pregnancy planned	Yes	50.1	46.5-53.8
	No	48.8	44.8-52.8
	Don't know	1.1	0.2-1.9

### *Infant HIV Exposure and MTCT Rate in Free State Province*

Text Box 2 shows that infants HIV exposure was 30.9% with a 1.2% early infant HIV infection prevalence and a 3.8% (95%CI: 2.3-5.3%) MTCT rate at 4-8 weeks is the highest in South Africa. The percentage of infants with reported HIV-negative mothers who were actually HIV-exposed (presumed maternal HIV acquisition) was 4.2% (95%CI: 3.0-5.3%) was significantly lower than in 2010.


**Text Box 2: Free State Infant HIV Exposure and MTCT**

2010			
Infant HIV Exposure % (95%CI)	Infant HIV infection prevalence at 4-8 weeks	MTCT @ 4-8 weeks: %(95%CI)	%Infants of reported HIV-negative mothers who had HIV antibody
31.3 (29.1-33.5)	2.4 (1.6-3.2)	5.9 (3.8-8.0)	5.4 (4.3-6.4)
2011			
Infant HIV Exposure % (95%CI)	Infant HIV infection prevalence at 4-8 weeks	MTCT @ 4-8 weeks: %(95%CI)	%Infants of reported HIV-negative mothers who had HIV antibody
30.9 (28.6-33.3)	1.2 (0.7-1.7)	3.8 (2.3-5.3)	4.2 (3.0-5.3)

### *PMTCT Service Uptake (PMTCT cascade) in the Free State Province*

Figure 7 shows uptake of HIV testing (97.64%) and coverage of ARV prophylaxis/ART is fairly high in Free State (96.05%). CD4 testing was at 63.15%. The high presumed maternal potential HIV acquisition may explain the high MTCT rate although, more detailed investigations are needed to examine 'effective coverage' of the PMTCT cascade (i.e., despite high CD4 cell and ART/prophylaxis uptake adherence to ARVs was low) and support for mothers post-HIV diagnosis and during pregnancy. Adherence and care and support data were not gathered in 2010.

**Figure 7 PMTCT service uptake (PMTCT cascade) in the Free State**



**Note:** The first three indicators apply to all mothers, while the last three apply only to those who reported being HIV-positive. Red indicates the percentage receiving ART while blue indicates the percentage receiving ARV prophylaxis.

### 3.6.3 Gauteng

The SAPMTCTE in Gauteng province attained 96% of targeted sample size.

#### *General Description of Provincial Sample*

Socioeconomic indicators show that compared to other provinces, participants in Gauteng Province have higher rates of flush toilet and electricity, gas or paraffin fuel source (Table 10).

**Table 10** Baseline characteristics of Gauteng SAPMTCTE survey participants

Characteristics	2010		
	Categories	%	95% CI
Relationship to child	Mother	98.4	97.8-99.0
	Caregiver	1.6	1.0-2.2
Mean age of mother (range)	26.6 (13-49)		
Infant gender	Male	52.1	49.8-54.3
	Female	47.9	45.7-50.2
Education of mother	None	1.5	0.9-2.2
	Grade 1-7	10.9	9.3-12.5
	Grade 8-12	80.2	77.7-82.7
	Above Grade 12	7.1	5.4-8.9
Marital status of mother	Single	69.9	65.3-74.4
	Married/cohabitating	29.4	24.8-33.9
Main building material of house	Brick/cement block	77.1	73.3-80.8
	Informal material	22.7	19.0-26.5
	Traditional material/mud	0.2	0.0-0.4
Main source of drinking water	Piped in house or yard	92.5	90.0-94.9
	Not piped in house or yard	7.5	5.1-9.9
Type of toilet	Flush toilet	84.8	81.6-88.0
	Pit latrine	14.5	11.5-17.6
	None	0.6	0.0-1.2
	Other	0.1	0.0-0.3
Main source of fuel	Electricity/gas/paraffin	99.2	98.8-99.6
	Other	0.8	0.4-1.2
Depletion of food supply in past 12 months	Yes	9.8	7.3-12.3
	No	89.9	87.4-92.4

2011			
Characteristics	Categories	%	95% CI
Relationship to child	Mother	98.4	98.0-99.9
	Father	0.2	0.0-0.4
	Grandmother/grandfather	0.9	0.5-1.3
	Guardian/legal guardian	0.2	0.0-0.5
	Caregiver	0.2	0.0-0.4
Mean age of mother (range)			
Infant gender	Male	51.0	48.9-53.0
	Female	49.0	47.0-51.1
Education of mother	None	1.2	0.8-1.7
	Grade 1-7	8.1	6.6-9.6
	Grade 8-12	81.6	78.7-84.5
	Completed tertiary/technical /university	9.0	6.8-11.1
	Don't know	0.1	0.0-0.3
Marital status of mother	Single	69.1	64.8-73.4
	Married	18.8	15.9-21.7
	Co-habiting	11.5	7.5-15.5
	Widowed	0.2	0.0-0.4
	Divorced/separated	0.3	0.0-0.6
	Don't know	0.1	0.0-0.2
Main building material of house	Brick/Cement block	80.5	76.6-84.4
	Informal material/corrugated iron/wood	19.5	15.6-23.4
	Traditional material/mud	Nil	
	other	Nil	
Main source of drinking water	Piped in house or yard	92.9	90.5-95.3
	Not piped in house or yard	7.1	4.7-9.5
Type of toilet	Flush toilet	87.1	83.5-90.6
	Pit latrine including ventilated pit latrine	12.6	9.1-16.2
	None	0.2	0.0-0.4
	Other	0.1	0.0-0.3
Main source of fuel	Electricity/gas/paraffin	99.3	98.7-99.8
	Other	0.8	0.2-1.3
Depletion of food supply in past 12 months	Yes	8.2	5.9-10.4
	No	91.7	89.5-94.0
	Don't know	0.13	0.0-0.3
Was this pregnancy planned	Yes	43.1	39.6-46.7
	No	56.8	53.2-60.3
	Don't know	0.1	0.0-0.3

### *Infant HIV Exposure and MTCT Rate in Gauteng*

Text Box 3 shows that infants' HIV exposure was 33.1%, with a 0.8% early infant HIV infection prevalence and a 2.1% (95%CI: 0.2-3.4%) MTCT rate at 4-8 weeks. Maternal potential HIV acquisition was 3.4% (95% CI 2.4-4.4%).


**Text Box 3: Gauteng Infant HIV Exposure and MTCT**

2010			
Infant HIV Exposure % (95%CI)	Infant HIV infection prevalence at 4-8 weeks	MTCT @ 4-8 weeks:(95%CI)	%Infants of reported HIV-negative mothers who had HIV antibody
30.4 (27.9-33.0)	1.1 (0.6-1.5)	2.5 (1.5-3.6)	3.0 (2.2-3.9)
2011			
Infant HIV Exposure % (95%CI)	Infant HIV infection prevalence at 4-8 weeks	MTCT @ 4-8 weeks:(95%CI)	%Infants of reported HIV-negative mothers who had HIV antibody
33.1 (29.8-36.4)	0.8 (0.3-1.2)	2.1 (0.2-3.4)	3.4 (2.4-4.4)

### *PMTCT service uptake (PMTCT cascade) in Gauteng*

Gauteng has achieved close to 100% (98.54% tested and 99.73% received test) HIV testing rate. Coverage of ARV prophylaxis or ART is 94.21% with 46.10% of women on ART, the highest of any province. However, CD4 count is only 77.06% and intended EID coverage while higher than other provinces is still below half (25.39%). (Figure 8)

**Figure 8 PMTCT Service Uptake (PMTCT cascade) in Gauteng**



**Note:** The first three indicators apply to all mothers, while the last three apply only to those who reported being HIV-positive. Red indicates the percentage receiving ART while blue indicates the percentage receiving ARV prophylaxis.



### 3.6.4 KwaZulu-Natal

The SAPMTCTE in KwaZulu-Natal attained 87% of targeted sample size.

#### *General Description of Provincial Sample*

KwaZulu-Natal had a very high rate of single mothers at 89.58%, while brick/cement block house was 55.34% and piped water were just above 65%, and 18.35% of participants described depletion of food supply in the past 12 months. (Table 11)

**Table 11 Baseline characteristics of KwaZulu-Natal SAPMTCTE survey participants**

2010			
Characteristics	Categories	%	95% CI
Relationship to child	Mother	95.5	95.4-97.5
	Caregiver	3.5	2.5-4.6
Mean age of mother (range)	24.9 (14-47)		
Infant gender	Male	47.9	45.2-50.7
	Female	52.1	49.3-54.8
Education of mother	None	1.5	0.7-2.3
	Grade 1-7	14.5	11.6-17.3
	Grade 8-12	79.5	76.3-82.6
	Above Grade 12	4.0	2.8-5.2
Marital status of mother	Single	90.7	89.0-92.4
	Married/cohabitating	8.9	7.2-10.6
Main building material of house	Brick/Cement block	61.9	55.5-68.3
	Informal material	13.3	9.0-17.6
	Traditional material/mud	24.8	18.3-31.4
Main source of drinking water	Piped in house or yard	60.6	52.8-68.4
	Not piped in house or yard	39.4	31.6-47.2
Type of toilet	Flush toilet	24.4	17.8-30.9
	Pit latrine	71.9	65.2-78.5
	None	3.8	1.0-6.5
	Other	0.0	--
Main source of fuel	Electricity/Gas/Paraffin	83.4	78.8-88.0
	Other	16.6	12.0-21.2
Depletion of food supply in past 12 months	Yes	21.6	16.2-27.0
	No	77.7	72.3-83.1

2011			
Characteristics	Categories	%	95% CI
Relationship to child	Mother	96.5	95.4-97.7
	Father	0.1	0.0-0.2
	Grandmother/grandfather	1.7	0.9-2.4
	Guardian/legal guardian	0.4	0.0-0.8
	Caregiver	1.4	0.6-2.1
Mean age of mother (range)			
Infant gender	Male	49.7	46.5-52.8
	Female	50.4	47.2-53.5
Education of mother	None	1.3	0.6-2.0
	Grade 1-7	13.8	10.8-16.7
	Grade 8-12	80.9	77.5-84.4
	Completed tertiary/technical /university	4.0	2.6-5.4
	Don't know	-	-
Marital status of mother	Single	89.6	85.7-93.4
	Married	7.3	5.6-9.0
	Co-habiting	3.1	0.0-6.8
	Widowed	-	-
	Divorced/separated	-	-
	Don't know	-	-
Main building material of house	Brick/Cement block	68.3	61.9-74.6
	Informal material/corrugated iron/wood	12.4	9.0-15.7
	Traditional material/mud	19.1	12.0-26.2
	other	0.2	0.0-0.4
Main source of drinking water	Piped in house or yard	55.3	46.3-64.3
	Not piped in house or yard	44.7	35.7-53.67
Type of toilet	Flush toilet	22.6	16.1-29.0
	Pit latrine including ventilated pit latrine	71.0	64.1-78.0
	None	6.4	1.4-11.4
	Other	-	-
Main source of fuel	Electricity/gas/paraffin	81.5	74.2-88.7
	Other	18.5	11.3-25.8
Depletion of food supply in past 12 months	Yes	18.3	13.5-23.2
	No	81.6	76.7-86.4
	Don't know	0.1	0.0-0.3
Was this pregnancy planned	Yes	22.7	19.3-26.0
	No	77.0	73.7-80.4
	Don't know	0.31	0.0-0.62

### *Infant HIV Exposure and MTCT Rate in KwaZulu-Natal*

Text Box 4 shows that infants HIV exposure was 44.4%, with a 0.9% early infant HIV infection prevalence and a 2.1% (95%CI: 0.9-3.3%) MTCT rate at 4-8 weeks. Among infants whose mothers reported being HIV negative 5.0% (95%CI: 3.7-6.4%) were HIV exposed (maternal potential HIV acquisition after the initial test), which was significantly higher than in 2010.


**Text Box 4: KwaZulu-Natal HIV Infant Exposure and MTCT**

2010			
Infant HIV Exposure % (95%CI)	Infant HIV infection prevalence at 4-8 weeks	MTCT @ 4-8 weeks: %(95%CI)	%Infants of reported HIV-negative mothers who had HIV antibody
44.3 (40.2-48.4)	1.9 (1.2-2.7)	2.9 (1.7-4.0)	3.2 (2.1-4.4)
2011			
Infant HIV Exposure % (95%CI)	Infant HIV infection prevalence at 4-8 weeks	MTCT @ 4-8 weeks: %(95%CI)	%Infants of reported HIV-negative mothers who had HIV antibody
44.4 (39.8-48.9)	0.9 (0.4-1.5)	2.1 (0.9-3.3)	5.0 (3.7-6.4)

### *PMTCT service uptake (PMTCT cascade) in the KwaZulu-Natal*

In KwaZulu-Natal, close to 100% of pregnant mothers receive testing and test results. More than 85% of HIV-positive mothers received their CD4 cell count test result. ARV/ART coverage was also high (95.55%). The high uptake of CD4 cell count results, together with the high percentage of pregnant women and infants receiving ARV prophylaxis or ART may explain the lower MTCT rate in KwaZulu-Natal. (Figure 9) Similar to other provinces, coverage of intended EID is low at 63.56%, but compared to other provinces, KwaZulu-Natal is the second highest in its EID coverage.

Figure 9 PMTCT service uptake (PMTCT cascade) in KwaZulu-Natal



**Note:** The first three indicators apply to all mothers, while the last three apply only to those who reported being HIV-positive. Red indicates the percentage receiving ART while blue indicates the percentage receiving ARV prophylaxis.

### 3.6.5 Limpopo

The SAPMTCT Evaluation in Limpopo province attained 73% of targeted sample size.

#### *General Description of Provincial Sample*

Limpopo showed a slightly higher rate of grandmother/grandfather (other than mothers) bringing infants to services (1.96%) and slightly lower percentage of single mothers (69.7%) compared to South African national rates. (Table 12)

**Table 12 Baseline characteristics of Limpopo SAPMTCTE survey participants**

2010			
Characteristics	Categories	%	95% CI
Relationship to child	Mother	93.8	92.4-95.2
	Caregiver	6.2	4.8-7.6
Mean age of mother (range)	26.0 (14-47)		
Infant gender	Male	50.3	47.4-53.2
	Female	49.7	46.8-52.6
Education of mother	None	1.6	1.0-2.3
	Grade 1-7	15.3	12.6-18.0
	Grade 8-12	75.0	71.9-78.1
	Above Grade 12	7.5	6.0-9.0
Marital status of mother	Single	69.7	66.0-73.4
	Married/cohabitating	30.0	26.4-33.7
Main building material of house	Brick/Cement block	89.2	87.2-91.2
	Informal material	8.3	6.3-10.2
	Traditional material/mud	2.5	1.6-3.5
Main source of drinking water	Piped in house or yard	47.4	41.5-53.4
	Not piped in house or yard	52.6	46.6-58.5
Type of toilet	Flush toilet	17.4	12.4-22.4
	Pit latrine	76.1	71.4-80.8
	None	6.0	4.2-7.9
	Other	0.4	0.1-0.8
Main source of fuel	Electricity/Gas/Paraffin	71.4	65.6-77.2
	Other	28.6	22.8-34.4
Depletion of food supply in past 12 months	Yes	15.1	12.0-18.1
	No	84.8	81.7-87.8

2011			
Characteristics	Categories	%	95% CI
Relationship to child	Mother	96.5	95.5-97.5
	Father	0.1	0.0-0.3
	Grandmother/grandfather	2.0	1.1-2.8
	Guardian/legal guardian	0.6	0.2-0.9
	Caregiver	0.8	0.3-1.4
Mean age of mother (range)			
Infant gender	Male	48.0	45.6-50.4
	Female	52.0	49.6-54.4
Education of mother	None	1.0	0.6-1.5
	Grade 1-7	11.6	9.4-13.7
	Grade 8-12	79.1	76.8-81.3
	Completed tertiary/technical /university	8.1	6.5-9.8
	Don't know	0.2	0.0-4
Marital status of mother	Single	62.7	58.7-66.7
	Married	32.1	28.2-35.9
	Co-habiting	5.0	2.8-7.1
	Widowed	-	-
	Divorced/separated	0.3	0.0-0.6
	Don't know	-	-
Main building material of house	Brick/Cement block	87.4	84.8-90.1
	Informal material/corrugated iron/wood	8.0	5.8-10.1
	Traditional material/mud	4.6	2.5-6.7
	other	-	-
Main source of drinking water	Piped in house or yard	42.9	37.9-47.9
	Not piped in house or yard	57.1	52.1-62.1
Type of toilet	Flush toilet	13.8	9.2-18.4
	Pit latrine including ventilated pit latrine	81.8	77.2-86.4
	None	4.3	2.7-5.9
	Other	0.1	0.0-0.3
Main source of fuel	Electricity/gas/paraffin	57.2	49.6-64.9
	Other	42.8	35.1-50.4
Depletion of food supply in past 12 months	Yes	17.1	14.0-20.3
	No	82.9	79.7-86.0
	Don't know	-	-
Was this pregnancy planned	Yes	52.6	49.9-55.3
	No	47.4	44.7-50.1
	Don't know	-	-

### Infant HIV Exposure and MTCT Rate in Limpopo

Text Box 5 shows that infants' HIV exposure was 23.0%, with a 0.8% early infant HIV infection prevalence and a 3.1% (95%CI: 1.2-4.9%) MTCT rate at 4-8 weeks. Among infants whose mothers reported being HIV-negative 1.6% (95%CI: 0.9-2.3) were HIV-exposed, which is a significant improvement over 2010.


**Text Box 5: Limpopo HIV Infant Exposure and MTCT**

2010			
Infant HIV Exposure % (95%CI)	Infant HIV infection prevalence at 4-8 weeks	MTCT @ 4-8 weeks:(95%CI)	%Infants of reported HIV-negative mothers who had HIV antibody
23.9 (21.8-25.9)	0.9 (0.4-1.5)	3.6 (1.4-5.8)	5.1 (3.6-6.7)
2011			
Infant HIV Exposure % (95%CI)	Infant HIV infection prevalence at 4-8 weeks	MTCT @ 4-8 weeks:(95%CI)	%Infants of reported HIV-negative mothers who had HIV antibody
23.0 (19.9-26.2)	0.8 (0.3-1.2)	3.1 (1.2-4.9)	1.6 (0.9-2.3)

### MTCT service uptake (PMTCT cascade) in the Limpopo

The PMTCT cascade shows that less than 70% of mothers received their CD4 cell count test result and 85.84% of HIV positive mothers received ARV prophylaxis or ART. (Figure 10)

Figure 10 PMTCT service uptake (PMTCT cascade) in Limpopo



**Note:** The first three indicators apply to all mothers, while the last three apply only to those who reported being HIV-positive. Red indicates the percentage receiving ART while blue indicates the percentage receiving ARV prophylaxis.

### 3.6.6 Mpumalanga

The SAPMTCTE attained 80% of targeted sample size in the province of Mpumalanga.

#### *General Description of Provincial Sample*

In Mpumalanga more than 80% of study participants have brick/cement block houses, piped water in house or yard, and electricity, paraffin or gas cooking fuel, but only 20.80% have a flush toilet.

Only 5.43% described depletion of food supply in the past 12 months. (Table 13)

**Table 13 Baseline characteristics of Mpumalanga SAPMTCTE survey participants**

2010			
Characteristics	Categories	%	95% CI
Relationship to child	Mother	95.0	94.3-95.8
	Caregiver	5.0	4.2-5.7
Mean age of mother (Range)	25.3 (13-46)		
Infant gender	Male	50.3	48.1-52.5
	Female	49.7	47.5-51.9
Education of mother	None	3.0	2.3-3.7
	Grade 1-7	17.9	15.7-20.2
	Grade 8-12	75.2	72.9-77.5
	Above Grade 12	2.1	1.3-2.9
Marital status of mother	Single	74.8	72.6-77.1
	Married/cohabitating	23.9	21.7-26.2
Main building material of house	Brick/Cement block	85.7	82.5-88.9
	Informal material	8.4	5.9-10.8
	Traditional material/mud	5.9	3.8-8.0
Main source of drinking water	Piped in house or yard	83.9	79.9-87.9
	Not piped in house or yard	16.1	12.1-20.1
Type of toilet	Flush toilet	30.3	23.9-36.7
	Pit latrine	66.5	60.1-72.8
	None	2.9	1.8-4.0
	Other	0.3	0.1-0.6
Main source of fuel	Electricity/Gas/Paraffin	88.3	65.6-77.2
	Other	11.7	22.8-34.4
Depletion of food supply in past 12 months	Yes	8.9	6.6-11.2
	No	89.1	86.6-91.6



2011			
Characteristics	Categories	%	95% CI
Relationship to child	Mother	97.6	97.0-98.3
	Father	0.3	0.0-0.6
	Grandmother/grandfather	1.0	0.5-1.4
	Guardian/legal guardian	0.5	0.2-0.9
	Caregiver	0.6	0.2-0.9
Mean age of mother (range)			
Infant gender	Male	50.4	48.5-52.2
	Female	49.6	47.8-51.5
Education of mother	None	2.5	1.6-3.4
	Grade 1-7	18.6	15.7-21.6
	Grade 8-12	76.5	73.5-79.6
	Completed tertiary/technical /university	2.1	1.4-2.8
	Don't know	0.2	0.0-0.5
Marital status of mother	Single	82.0	79.5-84.5
	Married	15.4	12.8-18.1
	Co-habiting	2.2	1.4-3.1
	Widowed	0.2	0.0-0.4
	Divorced/separated	0.1	0.0-0.3
	Don't know	-	-
Main building material of house	Brick/Cement block	89.9	87.2-92.5
	Informal material/corrugated iron/wood	8.0	5.5-10.4
	Traditional material/mud	2.1	0.9-3.3
	other	0.1	0.0-0.2
Main source of drinking water	Piped in house or yard	85.4	80.5-90.2
	Not piped in house or yard	14.6	9.8-19.5
Type of toilet	Flush toilet	20.8	14.9-26.7
	Pit latrine including ventilated pit latrine	78.9	73.0-84.8
	None	0.3	0.1-0.6
	Other	-	-
Main source of fuel	Electricity/gas/paraffin	97.3	96.0-98.6
	Other	2.7	1.3-4.0
Depletion of food supply in past 12 months	Yes	5.4	3.6-7.3
	No	94.4	92.5-96.4
	Don't know	0.1	0.0-0.3
Was this pregnancy planned	Yes	47.1	41.6-52.7
	No	52.8	47.3-58.3
	Don't know	0.1	0.0-0.2

### *Infant HIV Exposure and MTCT Rate in Mpumalanga*

Text Box 6 shows that infants' HIV-exposure was 35.60%, with a 1.2% early infant HIV infection prevalence and a 3.3% (95%CI: 2.2-4.5%) MTCT rate at 4-8 weeks. This is the second highest MTCT among South African provinces. Among infants whose mothers reported being HIV-negative 10.2% (95%CI: 8.2-12.2%) were HIV-exposed, the highest among South African provinces (along with Eastern Cape) and significantly worse than in 2010, suggesting major programmatic problems in Mpumalanga.


**Text Box 6: Mpumalanga HIV Infant Exposure and MTCT**

2010			
Infant HIV Exposure % (95%CI)	Infant HIV infection prevalence at 4-8 weeks	MTCT @ 4-8 weeks:(95%CI)	%Infants of reported HIV-negative mothers who had HIV antibody
37.0 (34.3-39.7)	3.0 (2.1-3.8)	5.7 (4.1-7.3)	7.8 (5.8-9.7)
2011			
Infant HIV Exposure % (95%CI)	Infant HIV infection prevalence at 4-8 weeks	MTCT @ 4-8 weeks:(95%CI)	%Infants of reported HIV-negative mothers who had HIV antibody
35.6 (33.3-37.8)	1.2 (0.8-1.7)	3.3 (2.2-4.5)	10.2 (8.2-12.2)

### *PMTCT service uptake (PMTCT cascade) in the Mpumalanga*

Mpumalanga also had a high coverage of testing. (Figure 11) However the high rate of HIV-exposed infants whose mothers reported being HIV-negative suggests that these mothers and babies would not have received any ARV prophylaxis and this, coupled with the poor PMTCT cascade above (<70% mothers receiving CD4 cell count results; 92.39% receiving ARV prophylaxis or ART and adherence to these regimens is not known i.e. 'effective coverage' is unknown) potentially explains the high MTCT rate in Mpumalanga province.

Figure 11 PMTCT service uptake (PMTCT cascade) in Mpumalanga



**Note:** The first three indicators apply to all mothers, while the last three apply only to those who reported being HIV-positive. Red indicates the percentage receiving ART while blue indicates the percentage receiving ARV prophylaxis.

### 3.6.7 Northern Cape

Northern Cape achieved a sample realisation of 63%. This was due to a high number of medium-size clinics in the sample requiring disproportionate resources.

#### *General Description of Provincial Sample*

In Northern Cape 80.44% of mothers were single. More than 80% of families had Electricity, piped water and flush toilets. Only 21.45% reported depletion of food supply in the past 12 months (Table 14)

**Table 14** Baseline characteristics of Northern Cape SAPMTCTE survey participants

Characteristics	2010		
	Categories	%	95% CI
Relationship to child	Mother	97.6	96.7-98.5
	Caregiver	2.4	1.5-3.3
Mean age (range)	25.8 (14-45)		
Infant gender	Male	48.9	46.1-51.7
	Female	51.1	48.3-53.9
Education of mother	None	3.5	2.6-4.4
	Grade 1-7	18.3	16.1-20.4
	Grade 8-12	74.3	71.8-76.9
	Above Grade 12	3.0	1.9-4.2
Marital status of mother	Single	78.0	75.0-81.1
	Married/cohabitating	21.1	18.0-24.1
Main building material of house	Brick/Cement block	80.4	77.5-83.4
	Informal material	18.7	15.9-21.5
	Traditional material/mud	0.9	0.4-1.3
Main source of drinking water	Piped in house or yard	93.5	91.6-95.4
	Not piped in house or yard	6.5	4.6-8.4
Type of toilet	Flush toilet	87.6	85.1-90.1
	Pit latrine	7.8	5.9-9.8
	None	2.2	1.4-3.0
	Other	2.4	1.7-3.0
Main source of fuel	Electricity/Gas/Paraffin	97.6	96.7-98.5
	Other	2.4	1.5-3.3
Depletion of food supply in past 12 months	Yes	10.9	8.1-13.6
	No	89.1	86.4-91.9

2011			
Characteristics	Categories	%	95% CI
Relationship to child	Mother	98.6	97.9-99.3
	Father	-	-
	Grandmother/grandfather	0.6	0.2-1.0
	Guardian/legal guardian	0.8	0.3-1.3
	Caregiver	-	-
Mean age of mother (range)			
Infant gender	Male	54.2	51.5-56.8
	Female	45.8	43.2-48.5
Education of mother	None	2.0	1.2-2.8
	Grade 1-7	14.0	11.7-16.4
	Grade 8-12	82.0	79.7-84.3
	Completed tertiary/technical /university	1.4	0.6-2.2
	Don't know	0.6	0.2-1.0
Marital status of mother	Single	80.4	77.5-83.4
	Married	18.0	14.8-21.1
	Co-habiting	1.0	0.1-1.9
	Widowed	0.6	0.2-1.0
	Divorced/separated	-	-
	Don't know	-	-
Main building material of house	Brick/Cement block	79.2	75.3-83.2
	Informal material/corrugated iron/wood	20.8	16.8-24.7
	Traditional material/mud	-	-
	other	-	-
Main source of drinking water	Piped in house or yard	90.3	87.6-93.0
	Not piped in house or yard	9.7	7.0-12.4
Type of toilet	Flush toilet	80.6	76.7-84.5
	Pit latrine including ventilated pit latrine	11.9	9.2-14.5
	None	3.4	2.0-4.7
	Other	4.2	2.5-5.8
Main source of fuel	Electricity/gas/paraffin	96.2	94.60-97.89
	Other	3.8	2.11-5.40
Depletion of food supply in past 12 months	Yes	21.4	18.5-23.8
	No	78.7	75.9-81.4
	Don't know	0.2	0.0-0.4
Was this pregnancy planned	Yes	38.7	34.5-42.9
	No	60.9	56.7-65.1
	Don't know	0.4	0.1-0.8

### *Infant HIV Exposure and MTCT Rate in the Northern Cape*

Text Box 7 shows that infants' HIV-exposure was 15.1%, with a 1.0% early infant HIV infection prevalence and a 6.1% (95%CI: 2.5-9.6%) MTCT rate at 4-8 weeks. The larger confidence interval attached to this estimate is due to the smaller sample size attained in Northern Cape.


**Text Box 7: Infant HIV Exposure and MTCT Rate Northern Cape**

2010			
Infant HIV Exposure % (95%CI)	Infant HIV infection prevalence at 4-8 weeks	MTCT @ 4-8 weeks:%(95%CI)	%Infants of reported HIV-negative mothers who had HIV antibody
16.0 (13.7-18.3)	0.3 (0.1-0.6)	1.4 (0.1-3.4)	2.2 (1.2-3.3)
2011			
Infant HIV Exposure % (95%CI)	Infant HIV infection prevalence at 4-8 weeks	MTCT @ 4-8 weeks:%(95%CI)	%Infants of reported HIV-negative mothers who had HIV antibody
15.1 (12.7-17.5)	1.0 (0.4-1.6)	6.1 (2.5-9.6)	1.9 (1.2-2.7)

### *PMTCT service uptake (PMTCT cascade) in the Northern Cape*

Northern Cape has the lowest intended early infant diagnosis coverage of all provinces. Very few (11.59%) HI- positive mothers indicated an intention to receive infant testing service at the 6-week immunisation visit. However, that CD4-cell count uptake is reported in the Northern Cape, at 76.81%. Other PMTCT cascade indicators of the province have above 90% coverage, similar to other provinces. The rate of maternal potential HIV acquisition was 0.48% (upper limit 0.91%). The low MTCT rate is likely due to the high coverage of the PMTCT cascade in Northern Cape and the lower HIV acquisition rate compared with other provinces.

**Figure 12 PMTCT service uptake (PMTCT cascade) in the Northern Cape**



**Note:** The first three indicators apply to all mothers, while the last three apply only to those who reported being HIV-positive. Red indicates the percentage receiving ART while blue indicates the percentage receiving ARV prophylaxis.

### 3.6.8 The North West Province

The SAPMTCTE in the North West province attained 98% of targeted sample size.

#### General Description of Provincial Sample

In North West province 81.06% of mothers were single. Just over 70% of families had brick/cement houses and piped water, but fewer than half had a flush toilet (43.58%). Reported depletion of food supply in the last 12 months was 10.81%. (Table 15)

**Table 15 Baseline characteristics of North West SAPMTCTE survey participants**

2010			
Characteristics	Categories	%	95% CI
Relationship to child	Mother	97.0	96.1-97.9
	Caregiver	3.0	2.1-3.9
Mean age (range)	26.3 (14-46)		
Infant gender	Male	50.9	48.9-52.8
	Female	49.1	47.2-51.1
Education of mother	None	5.3	3.6-7.0
	Grade 1-7	19.1	16.1-22.0
	Grade 8-12	71.8	67.9-75.8
	Above Grade 12	3.3	2.2-4.4
Marital status of mother	Single	83.1	80.8-85.6
	Married/cohabitating	16.0	13.6-18.4
Main building material of house	Brick/Cement block	73.8	70.7-76.9
	Informal material	24.0	21.1-26.9
	Traditional material/mud	2.2	1.2-3.1
Main source of drinking water	Piped in house or yard	75.9	71.3-80.4
	Not piped in house or yard	24.1	19.6-28.7
Type of toilet	Flush toilet	44.1	37.1-51.0
	Pit latrine	54.0	47.1-61.0
	None	1.3	0.8-1.9
	Other	0.6	0.1-1.0
Main source of fuel	Electricity/Gas/Paraffin	93.5	91.4-95.5
	Other	6.5	4.5-8.6
Depletion of food supply in past 12 months	Yes	19.1	16.4-21.7
	No	80.6	78.0-83.3



2011			
Characteristics	Categories	%	95% CI
Relationship to child	Mother	97.4	96.6-98.1
	Father	0.2	0.0-0.4
	Grandmother/grandfather	1.8	1.2-2.4
	Guardian/legal guardian	0.3	0.1-0.5
	Caregiver	0.4	0.1-0.7
Mean age of mother (range)			
Infant gender	Male	46.8	44.5-49.2
	Female	53.2	50.8-55.5
Education of mother	None	3.5	2.4-4.5
	Grade 1-7	21.0	18.5-23.6
	Grade 8-12	70.9	67.8-73.9
	Completed tertiary/technical /university	4.5	3.2-5.8
	Don't know	0.2	0.0-0.3
Marital status of mother	Single	81.1	78.4-83.8
	Married	16.6	14.1-19.0
	Co-habiting	1.8	1.1-2.4
	Widowed	0.4	0.1-0.8
	Divorced/separated	0.2	0.0-0.4
	Don't know	-	-
Main building material of house	Brick/Cement block	74.5	71.2-77.8
	Informal material/corrugated iron/wood	23.5	20.2-26.7
	Traditional material/mud	2.0	1.0-3.2
	other	-	-
Main source of drinking water	Piped in house or yard	74.8	68.9-80.7
	Not piped in house or yard	25.2	19.3-31.1
Type of toilet	Flush toilet	43.6	35.8-51.4
	Pit latrine including ventilated pit latrine	54.3	46.6-62.0
	None	2.0	1.3-2.8
	Other	0.1	0.0-0.2
Main source of fuel	Electricity/gas/paraffin	96.0	94.8-97.2
	Other	4.0	2.8-5.2
Depletion of food supply in past 12 months	Yes	10.8	8.5-13.1
	No	88.5	86.29-90.80
	Don't know	0.7	0.26-1.03
Was this pregnancy planned	Yes	34.5	31.44-37.48
	No	64.7	61.68-67.77
	Don't know	0.8	0.27-1.37

### *Infant HIV Exposure and MTCT Rate in North West Province*

Text Box 8 shows that infants' HIV-exposure was 30.8%, with a 0.8% early infant HIV infection prevalence and a 2.6 (95% CI 1.1-4.0%) MTCT rate at 4-8 weeks. The percentage of infants with reported HIV-negative mothers who were actually HIV-exposed (presumed maternal HIV acquisition) was 3.7% (95%CI: 2.5-4.9%) was significantly lower than in 2010.


Text Box 8: Infant HIV Exposure and MTCT Rate in North West Province

2010			
Infant HIV Exposure % (95%CI)	Infant HIV infection prevalence at 4-8 weeks	MTCT @ 4-8 weeks: %(95%CI)	% Infants of reported HIV-negative mothers who had HIV antibody
31.3 (29.0-33.5)	1.9 (1.2-2.5)	4.4 (2.9-5.9)	5.4 (3.9-6.8)
2011			
Infant HIV Exposure % (95%CI)	Infant HIV infection prevalence at 4-8 weeks	MTCT @ 4-8 weeks: %(95%CI)	% Infants of reported HIV-negative mothers who had HIV antibody
30.8 (28.5-33.1)	0.8 (0.4-1.2)	2.6 (1.1-4.0)	3.7 (2.5-4.9)

### *PMTCT Service Uptake (PMTCT cascade) in the North West Province*

North West study participants had close to 100% HIV testing rate, but lower coverage of CD4 count (74.16%) and EID services. The province ranked as the second lowest in intended EID coverage. Only 12.99% of HIV-positive mothers in the sample intended to receive EID at the time of immunisation visit. The higher rate of infants with reported HIV-negative mothers who were HIV-exposed and who would not have received any PMTCT interventions, coupled with a only a 93.29% coverage of ARV prophylaxis / ART may explain the somewhat higher MTCT rate measured in the North West province.

**Figure 13 PMTCT service uptake (PMTCT cascade) in the North West Province**



**Note:** The first three indicators apply to all mothers, while the last three apply only to those who reported being HIV-positive. Red indicates the percentage receiving ART while blue indicates the percentage receiving ARV prophylaxis.

### 3.6.9 Western Cape

The SAPMTCTE in the Western Cape attained 99% of targeted sample size.

#### *General Description of Provincial Sample*

The majority of Western Cape participants reported use of piped water (in house) (98.01%), flush toilet (92.61%) and electricity, gas or paraffin (99.56%) for their fuel needs. However, a substantial percentage (29.74%) of the participants reported living in a house built from informal materials. A large percentage (14.80%) of respondents also reported that they experienced food shortage at least once in the last 12 months. (Table 16)

**Table 16** Baseline characteristics of Western Cape SAPMTCTE survey participants

Characteristics	Categories	%	95% CI
Relationship to child	Mother	97.1	96.3-97.9
	Caregiver	2.9	2.1-3.7
Mean age (range)	26.4 (14-47)		
Infant gender	Male	49.7	47.4-52.0
	Female	50.3	48.0-52.6
Education of mother	None	0.8	0.3-1.3
	Grade 1-7	15.1	12.2-17.9
	Grade 8-12	76.2	73.0-79.4
	Above Grade 12	7.3	5.1-9.5
Marital status of mother	Single	54.3	50.5-58.1
	Married/cohabitating	44.0	40.2-47.8
Main building material of house	Brick/Cement block	68.5	63.3-73.7
	Informal material	31.1	26.0-36.3
	Traditional material/mud	0.4	0.1-0.6
Main source of drinking water	Piped in house or yard	93.9	91.9-95.8
	Not piped in house or yard	6.1	4.2-8.0
Type of toilet	Flush toilet	90.7	88.4-93.0
	Pit latrine	5.6	3.5-7.7
	None	1.3	0.6-2.0
	Other	2.4	1.5-3.3
Main source of fuel	Electricity/Gas/Paraffin	99.3	99.0-99.7
	Other	0.7	0.3-1.0
Depletion of food supply in past 12 months	Yes	26.1	22.9-29.2
	No	73.0	69.9-76.0

2011			
Characteristics	Categories	%	95% CI
Relationship to child	Mother	97.29	96.58-98.01
	Father	0.21	0.01-0.40
	Grandmother/grandfather	1.01	0.53-1.49
	Guardian/legal guardian	0.29	0.06-0.51
	Caregiver	1.21	0.67-1.75
Mean age of mother (range)			
Infant gender	Male	49.35	47.33-51.37
	Female	50.65	48.63-52.67
Education of mother	None	0.69	0.23-1.15
	Grade 1-7	11.00	8.89-13.11
	Grade 8-12	81.55	78.98-84.12
	Completed tertiary/technical /university	6.32	4.42-8.21
	Don't know	0.45	0.16-0.74
Marital status of mother	Single	62.30	57.82-66.77
	Married	29.23	25.61-32.85
	Co-habiting	7.27	4.95-9.60
	Widowed	0.22	0.02-0.42
	Divorced/separated	0.98	0.60-1.37
	Don't know	-	-
Main building material of house	Brick/Cement block	70.26	64.11-76.42
	Informal material/corrugated iron/wood	29.74	23.59-35.90
	Traditional material/mud	-	-
	other	-	-
Main source of drinking water	Piped in house or yard	98.01	96.90-99.12
	Not piped in house or yard	1.99	0.88-3.10
Type of toilet	Flush toilet	92.61	90.01-95.22
	Pit latrine including ventilated pit latrine	5.26	2.83-7.69
	None	1.20	0.57-1.83
	Other	0.93	0.46-1.40
Main source of fuel	Electricity/gas/paraffin	99.56	99.28-99.84
	Other	0.44	0.16-0.72
Depletion of food supply in past 12 months	Yes	14.80	12.15-17.45
	No	84.72	82.04-87.40
	Don't know	0.48	0.17-0.79
Was this pregnancy planned	Yes	38.36	35.76-40.95
	No	61.64	59.05-64.24
	Don't know	-	-

### *Infant HIV Exposure and MTCT Rate in Western Cape*

Text Box 9 shows that infants' HIV-exposure was 17.8%, with a 0.4% early infant HIV infection prevalence and a 1.98% (95% CI 0.65-3.31%) MTCT rate at 4-8 weeks, which was significantly lower than in 2010. The percentage of infants with reported HIV-negative mothers who were actually HIV-exposed (presumed maternal HIV acquisition) was 0.7% (95%CI: 0.3-1.2%) was the lowest in South Africa. These data potentially suggest the Western Cape has the best quality and most effective PMTCT programme.


Text Box 9: Infant HIV Exposure and MTCT Rate in Western Cape

2010			
Infant HIV Exposure % (95%CI)	Infant HIV infection prevalence at 4-8 weeks	MTCT @ 4-8 weeks: %(95%CI)	%Infants of reported HIV-negative mothers who had HIV antibody
21.0 (30.7-33.3)	0.9 (0.4-1.5)	3.9 (1.9-5.8)	1.1 (0.3-1.9)
2011			
Infant HIV Exposure % (95%CI)	Infant HIV infection prevalence at 4-8 weeks	MTCT @ 4-8 weeks: %(95%CI)	%Infants of reported HIV-negative mothers who had HIV antibody
17.8 (14.8-20.8)	0.4 (0.1-0.6)	1.98 (0.65-3.31)	0.7 (0.3-1.2)

### *PMTCT service uptake (PMTCT cascade) in the Western Cape*

The Western Cape had close to 100% (97.73%) antenatal HIV testing rate. Almost all (99.41%) mothers who had antenatal HIV testing received their result. About 97.41% of the HIV-positive mothers were on either ART or ARV prophylaxis (including infant prophylaxis if mother is not on ART). Similar to other provinces, the majority of the mothers did not plan to receive EID services during the six-week immunisation visit.

**Figure 14 PMTCT service uptake (PMTCT cascade) in the Western Cape Province**



**Note:** The first three indicators apply to all mothers, while the last three apply only to those who reported being HIV-positive. Red indicates the percentage receiving ART while blue indicates the percentage receiving ARV prophylaxis.

### 3.7 Infant Feeding

HIV-positive mothers who recalled receiving infant feeding counselling during antenatal care was 89.2% with a range from 77.9% in Limpopo to 92.4% in Gauteng and KwaZulu-Natal.

Among all infants (regardless of HIV exposure status) 44.8% (95% CI 43.2-46.3%) were mixed breastfeeding; 28.0% (95% CI 25.6-29.4%) were exclusively breastfed in the 8 days prior to the 4-8 week interview and 27.2% (95% CI 26.1-28.3%) received no breast milk.

We categorised HIV-exposed infants who received breast milk plus any other milk or food (not including prescribed medicines) over the past eight days as being at-risk as they were practicing mixed breastfeeding. This ranged from a low of 9.6% in Gauteng to a high of 33.0% in Limpopo, with a national average of 14.0% (Table 17). It is interesting to note that the province with the lowest rate of infant feeding counselling (82% in Limpopo province) had high rates of at risk infant feeding in HIV-exposed infants (33%).



**Table 17 Infant feeding practices amongst HIV exposed infants over the past 8 days by province**

Province	Mother reportedly received infant feeding counseling during ANC % (95% CI)		At Risk/ Mixed Feeding % (95% CI)		Exclusive Breastfeeding	
	2010	2011	2010	2011	2010	2011
Eastern Cape	82.4 (76.1-88.8)	94.2 (91.3-97.3)	20.3 (16.6-23.9)	9.9 (6.7-13.0)	14.8 (11.3-18.2)	23.2 (17.8-28.6)
Free State	91.6 (88.7-94.6)	95.5 (93.7-97.4)	22.9 (19.1-26.8)	19.2 (14.9-23.5)	18.0 (14.5-21.5)	35.1 (30.1-40.1)
Gauteng	92.4 (89.7-95.1)	91.8 (87.9-95.6)	14.8 (11.4-18.1)	9.6 (6.2-13.0)	19.6 (15.9-23.2)	37.6 (32.2-43.0)
KwaZulu-Natal	92.4 (90.0-94.8)	97.2 (95.2-99.2)	14.0 (10.5-17.5)	10.4 (6.9-13.9)	26.1 (21.2-31.1)	42.5 (37.1-47.9)
Limpopo	77.9 (72.0-83.7)	82.2 (76.2-88.2)	32.8 (27.0-38.7)	33.0 (25.9-40.1)	20.3 (15.5-25.1)	28.3 (21.5-35.0)
Mpumalanga	91.5 (88.4-94.6)	93.5 (91.5-95.4)	29.7 (26.0-33.4)	20.6 (15.5-25.6)	13.9 (11.01-16.8)	34.5 (28.6-40.5)
Northern Cape	81.0 (73.3-88.6)	94.2 (90.3-98.2)	23.9 (16.1-31.8)	21.7 (13.8-29.7)	43.7 (37.0-50.3)	43.5 (35.9-51.1)
North West	81.5 (76.1-86.8)	90.6 (87.3-94.0)	21.3 (17.5-25.1)	23.2 (16.7-29.7)	25.7 (21.3-30.0)	38.9 (33.0-44.8)
Western Cape	85.3 (80.5-90.1)	95.0 (92.6-97.3)	7.0 (4.3-9.6)	11.4 (7.3-15.4)	7.9 (5.3-10.4)	18.8 (13.9)
South Africa	89.2 (87.8-90.6)	93.3 (92.0-94.7)	18.1 (16.5-19.7)	14.0 (12.3-15.7)	20.4 (18.5-22.3)	35.5 (33.1-38.0)

There are several encouraging findings with regards to infant feeding:

- Among mothers of HIV-exposed infants:
  - 35.5% (33.12-38.0%) reported exclusive breastfeeding over the past 8 days, which is a significant increase from the 20.4% (95% CI 18.5-22.3%) reported in 2010.
  - The prevalence of exclusive breastfeeding measured in the 2011 survey was lowest in the Western Cape Province and highest in Kwa Zulu Natal and Northern Cape Provinces. However in at least all provinces except for Limpopo and Northern Cape a significant increase in exclusive breastfeeding was measured between 2010 and 2011
  - 47.1% (44.9-49.3%) reported avoiding breastmilk, which is a reduction from the 61.5% (59.2-63.8) measured in 2010.
  - There was a significant increase in reported infant feeding counseling between 2010 and 2011 and the significant reduction in mixed feeding (at risk feeding) between 2010 and 2011.
  
- Amongst mothers of HIV unexposed infants:
  - 43.6% (41.6%-45.7%) reported exclusive breastfeeding over the past 8 days in 2011 compared with 31.3% (29.-33.0%) in 2012, illustrating a significant increase in exclusive breastfeeding in 2011 compared with 2010.
  - The Eastern Cape and Limpopo provinces recorded the lowest prevalence of exclusive breastfeeding. This is especially worrying as these are both rural provinces. The prevalence of exclusive breastfeeding was highest in the Kwa Zulu Natal and Mpumalanga provinces.
  - The prevalence of exclusive breastfeeding showed a significant increase between the 2010 and 2011 surveys in all provinces except Limpopo, Northern Cape and North West provinces.
  - Data also show a reduction in mixed feeding amongst HIV unexposed infants: In 2010 57.4% (55.5-59.2%) reported mixed feeding whereas in 2011 this dropped to 46.2% (44.2-48.3%).


## 4. DISCUSSION

### 4.1 Infant HIV Exposure

Figure 15 indicates the 2011 maternal HIV prevalence in mothers from the antenatal sentinel surveillance by province (NDOH, 2012).<sup>14</sup>

The weighted proportion of mothers who reported being HIV positive in the 2011 SAPMTCT Evaluation was 29.6% (95% CI 28.0-31.1%) which is very similar to the antenatal HIV prevalence which was measured as 29.5% (28.7-30.2%) in the 2011 national antenatal survey. The infant HIV exposure was higher, viz. weighted proportion of children who were HIV exposed at six weeks (4-8 weeks post-delivery) in the SAPMTCTE was 32.2% (95% confidence interval of 30.7-33.6%) It is expected for the SAPMTCTE infant HIV exposure to be result is slightly higher than the antenatal HIV prevalence as it measures incident HIV infections and sero-conversions.

Figure 15 Maternal antenatal HIV prevalence by province in South Africa




### 4.2 Mother-to-Child Transmission of HIV

Figure 16 presents data from the National Health Laboratory Service (NHLS) data warehouse for infants less than 2 months old.<sup>10</sup> The data show an increase in the number of HIV PCR tests done between 2003 and 2011; by 2011 58% of all HIV PCR tests were done early in infants. According to routine NHLS data early vertical HIV transmission in children <2 months of age dropped from 9.7% in 2008 to 2.8% in 2011. The NHLS data mirror the SAPMTCTE findings, and illustrate the gains made in

preventing early infants HIV infections. While these different data sources have varying methods, limitations and strengths the underlying message is that MTCT has reduced substantially in South Africa over the past three years and at present is less than 5%.

Figure 16 NHLS Early Infant Diagnosis PCR <2 months old 2011 (Sherman, IAS 2012)<sup>10</sup>



The MTCT rate measured at six weeks in recent PMTCT studies (not operational settings) using somewhat similar regimens to the South African 2010 PMTCT policy include:

- SWEN (2008) - Sd NVP peripartum and daily in infant until 6 weeks; 2.5% MTCT at 6 weeks;<sup>15</sup>
- Mitra (Kilewo et.al., 2008) - ZDV/3TC peripartum and 3TC postnatally; MTCT 3.8% at 6 weeks);<sup>16</sup> and
- Kesho Bora (2011) - AZT/NVP in mothers vs. ART for mothers; MTCT 5.0% versus 3.3% respectively at 6 weeks)<sup>17</sup>.

The SAPMTCTE results compare favourably with these results. Achievement of results similar to trials in a national PMTCT programme is very encouraging.

### **Provincial Variation in MTCT**

There was a greater than 4-fold difference in MTCT across the 9 provinces in South Africa. The provincial variation in MTCT is due to the differences in 'effective coverage' and quality of the PMTCT programme including uptake of CD4 cell count testing results, repeat HIV testing at 32 weeks, appropriate ARV prophylaxis/ART for HIV-positive women, and adherence to PMTCT regimens. More detailed explorations of quality and adherence to PMTCT prophylaxis or ART are underway to understand MTCT rates across provinces.

### 4.3 PMTCT Cascade

Missed opportunities along the PMTCT cascade of services (Stringer et.al., 2003) can reduce both the coverage and quality of the PMTCT programme. HIV Testing in ANC clinics is the entry point into the PMTCT programme. High coverage of this and each subsequent step reduces missed opportunities for care. In 2011, ANC HIV testing by mothers was almost universal 98.3% but services further along the cascade were not as high, with only 77.4% of HIV-positive mothers getting a CD4 count. These were similar to 2010 data. The percentage of mothers and infants received ARV prophylaxis or treatment according to protocol increased slightly to 93.9% (from 91.3% in 2010). This data does show improvement over a previous report from KwaZulu-Natal where prior to a quality improvement intervention only 85% of women were tested in ANC, 40% received a CD4 test and only 15% were given appropriate ARV prophylaxis (Doherty et.al., 2009). After the intervention, the data from the Doherty study was comparable to the SAPMTCTE with 98% ANC HIV testing, 97% CD4 testing and 68% appropriate ARV prophylaxis. There has been an effort in South Africa in the last few years to improve the PMTCT programme through interventions like the one described by Doherty et.al. (2009) as well as others (e.g. Best Practices in Prevention of Mother-to-Child Transmission (PMTCT) of HIV South Africa; NDOH/MRC/UWC/UNICEF/USAID, 2009). These efforts are clearly impacting PMTCT, as shown by programme indicators and infant outcomes (early MTCT) as described in this report.

Of ALL mothers enrolled in the survey, 29.4% reported being HIV-positive while HIV antibody was found in 32% of ALL infants. Of concern is that of those *mothers who reported being HIV-negative*, 3.9% of their infants had HIV antibodies, suggesting a high rate of maternal potential acquisition of HIV infection during pregnancy. This rate also varied substantially across provinces from a low of 0.7% in the Western Cape to a high of 10.2% in Mpumalanga. The indicator 'Maternal potential HIV acquisition' is a combination of the following scenarios:

- i. Mothers do not wish to admit being HIV-positive and reported being HIV negative. The data show that refusals for infant HIV testing were low and disclosure was high; thus the contribution that this scenario makes to the indicator is probably minimal.
- ii. Mothers were tested during the window period for the ANC test.
- iii. Poor QC/performance of rapid tests in the field caused false negative results at ANC on HIV-infected women. Reported field sensitivities are as low as 87% to 95% depending on the rapid test. In correlation to this was mothers who reported being HIV-positive but for which infant test HIV antibody negative which was 2.5% (95% CI 1.8-3.2%) and also suggests potential problems with performance of rapid tests during ANC.
- iv. True acquisition of HIV after the last HIV test – which for most mothers was during pregnancy.

Regardless of the cause this group of women and infants represent a substantial missed opportunity for care as the mothers and infants did not receive ARV prophylaxis or appropriate counselling and represents a metric for PMTCT programme quality. The high rate of maternal potential HIV

acquisition is also worrying as it is higher than previously reported (3%) in a study conducted in KwaZulu-Natal (Moodley et al., 2009).

#### **4.4 Early Infant Diagnosis**

Intention to obtain a PCR test at the six week immunization visit remain the same nationally between 2010 and 2011 [35.1% (30.6-39.6%) versus 38.5% (34.3-42.6%)] respectively illustrating persistent gaps in the integration between routine child health services and HIV-related care. The Free State province is particularly concerning as 2011 data show that intention for infant HIV testing at six weeks immunisation visits significantly reduced between 2010 and 2011 [43.7% (33.3-54.1%) versus 24.6% (17.8-31.4%)] respectively. Marginally significant improvements in integration are seen from 2010 to 2011 in the Western Cape [37.9% (28.8-47.0%) versus 46.3% (36.1-56.5%)] and Mpumalanga provinces [29.8% (23.1—36.5%) versus 41.2% (32.1-50.3%)] respectively and significant increases in integration are seen in KZN province [41.1% (30.5-51.5%) in 2010 versus 63.6% (53.6-73.5%) in 2011].

#### **4.5 Infant Feeding**

Infant feeding per caregiver recall for the past 8 days suggests a substantial increase in infant feeding counseling, increase in exclusive breastfeeding and reduction in mixed breastfeeding amongst HIV exposed and unexposed infants. The Tshwane Declaration of Support for Breastfeeding was adopted in August 2011 just as the survey started.<sup>6</sup> It received wide publicity and its effects are likely to have been measured over the duration of the survey.

## 5. STRENGTHS AND LIMITATIONS OF SAPMTCTE

### Strengths

This evaluation provides estimates of early vertical HIV transmission using a national and provincial population-based representative sample of infants 4-8 weeks of age. The survey was conducted 16 months after South Africa adopted PMTCT Option A, and thus provides population level data on effectiveness of WHO PMTCT Option A. It also provides data on uptake of the PMTCT programme and infant feeding.

### Limitations

The survey has the following limitations:

The data is facility-based using infants presenting for immunisation. Infants who do not come for immunisation or have already died by 6 weeks of age are not included in the sample suggesting a possible under-estimation of infant HIV infection prevalence.

Maternal Incidence (sero-conversion during pregnancy) was based on self-reports of previous HIV-negative status and presence of HIV antibodies in infant ELISA test. Mothers may not accurately report their previous HIV status for a variety of reasons, such as fear of stigma and disclosure. Confidentiality was assured and discussed as part of the informed consent process and a private place was secured for the conduct of interviews in an attempt to reduce this potential limitation.

Coverage of PMTCT programme indicators was via maternal recall and was not verified with maternal antenatal or intrapartum records, however the recall period was relatively short (generally less than 3-6 months).

Two-stage cluster random sampling was used. The primary sampling unit was primary health care clinics reporting at least 130 immunisations per year from the 2007 DHIS data. Therefore this sample excluded smaller primary health care facilities due to logistic reasons and secondary and tertiary facilities, mobile clinics and other facilities in order to focus on PMTCT in the primary health care services. Therefore this survey is not representative of these excluded facilities.

Finally, this survey does not measure postnatal HIV transmission.

## 6. CONCLUSION AND RECOMMENDATIONS

---

### Conclusions:

1. South Africa reduced early vertical HIV transmission from 3.5% in 2010 to 2.7% (95% CI 2.1-3.2%) in 2011. This followed the adoption of WHO PMTCT Option A.
2. Reported infant feeding counseling and infant feeding practices improved between 2010 and 2011.
3. Uptake of CD4 cell count testing and results is still poor (78% of HIV positive women)

### Recommendations:

1. Bottle necks to further reductions in MTCT need to be explored
2. The effectiveness of PMTCT Option B needs to be measured: will it result in further reductions in vertical HIV transmission as the need for CD4 cell count testing to initiate antiretroviral treatment is removed
3. The long-term effectiveness of PMTCT needs to be urgently measured
4. Integration between routine child health services and early infant diagnosis needs to be intensified
5. More work is needed to provide appropriate infant feeding counseling for continued improvements in infant feeding practices



## REFERENCES

---

1. National Department of Health. Policy and Guidelines for the Implementation of the PMTCT programme. 2008. Available from <http://www.info.gov.za/view/DownloadFileAction?id=77877>. Accessed 1 October 2009.
2. National Department of Health and SANAC. Clinical Guidelines: PMTCT (Prevention of Mother-to-Child Transmission). 2010. Available from [http://www.fidssa.co.za/images/PMTCT\\_Guidelines.pdf](http://www.fidssa.co.za/images/PMTCT_Guidelines.pdf). Accessed 6 January 2014.
3. Goga A, Dinh T, Jackson D, for the SAPMTCTE Study Group. Evaluation of the Effectiveness of the National Prevention of Mother-to-Child Transmission (PMTCT) Programme Measured at Six Weeks Postpartum in South Africa, 2010. Published 2012. Available from <http://www.mrc.ac.za/healthsystems/reports.htm>. Accessed 6 January 2014
4. National Department of Health SA. The National Strategic Plan on HIV, STIs and TB, 2012-2016. Available from [www.info.gov.za/view/DownloadFileAction?id=155622](http://www.info.gov.za/view/DownloadFileAction?id=155622). Accessed 18 October 2013.
5. National Department of Health SA. National Department of Health. Protocol for providing a comprehensive package of care for the prevention of mother to child transmission of HIV (PMTCT) in South Africa. 2001. Available from [http://www.law-lib.utoronto.ca/Diana/TAC\\_case\\_study/Protocol.html](http://www.law-lib.utoronto.ca/Diana/TAC_case_study/Protocol.html). Accessed 18 October 2013.
6. National Department of Health Breastfeeding Summit. The Tshwane Declaration of Support for breastfeeding in South Africa. *South African Journal of Clinical Nutrition* 2011; 24(4): 214-6.
7. Rollins N, Little K, Mzoloa S, Horwood C, Newell M-L. Surveillance of mother-to-child transmission prevention programmes at immunization clinics: the case for universal screening. *AIDS* 2007; 21: 1341-7.
8. Rollins N, Mzolo S, Moodley T, Esterhuizen T, van Rooyen H. Universal HV testing of infants at immunisation clinics: an acceptable and feasible approach for early infant diagnosis in high HIV prevalence settings. *AIDS* 2009; 23(1851-1857).
9. Chantry C, Cooper E, Pelton S, Zorilla C, Hillyer G, Diaz C. Seroreversion in human immunodeficiency virus-exposed but uninfected infants. *Pediatr Infect Dis J* 1995; 14(5): 382-7.
10. Sherman G, Lilian R, Coovadia A. The performance of five rapid HIV tests using whole blood in infants and children – selecting a test to achieve the clinical objective. *Paed Infect Dis J* 2012; 31(3): 267-72.
11. National Department of Health. 2008 National Antenatal Sentinel HIV and Syphilis Prevalence Survey, South Africa. Published 2009. Available from <http://www.info.gov.za/view/DownloadFileAction?id=109007>. Accessed 18 October 2013.
12. Lehtonen R, Pahkinen E. *Practical Methods for Design and Analysis of Complex Surveys* 2nd ed. West Sussex, UK: John Wiley & Sons; 2004.
13. Stevens W, Erasmus L, Moloji M, Taleng T, Sarang S. Performance of a novel human immunodeficiency virus (HIV) type 1 total nucleic acid-based real-time PCR assay using whole blood and dried blood spots for diagnosis of HIV in infants. *J Clin Microbiol* 2008; 46: 3941-5.
14. National Department of Health. The 2011 National Antenatal Sentinel HIV & Syphilis Prevalence Survey in South Africa. Pretoria, Published 2012. Available from <http://health-e.org.za/documents/f0980fb5107a7ce543a8bd5730e52333.pdf>. Accessed 18 October 2013.
15. Six week extended dose nevirapine (SWEN) Study team et al. Extended dose nevirapine at 6 weeks of age for infants to prevent HIV transmission via breastfeeding in Ethiopia, India and Uganda: an analysis of 3 randomised controlled trials. *The Lancet* 2008; 372: 300-13.

16. Kilewo C, Karlsson K, Massawe A, et al. Prevention of Mother-to-Child Transmission of HIV-1 through breastfeeding by treating infants prophylactically with lamivudine in Dar es Salaam. *JAIDS* 2008; 48(3): 315-23.
17. Kesho Bora Study Group, de Vincenzi I. Triple antiretroviral compared with zidovudine and single-dose nevirapine prophylaxis during pregnancy and breastfeeding for prevention of mother-to-child transmission of HIV-1 (Kesho Bora study): a randomised controlled trial. *Lancet Infect Dis*; 2011; Mar;11(3): 171-80.

## APPENDIX I: SAMPLE SIZE AND SAMPLING

(See next page)

## SAMPLE SIZE CALCULATION BY PROVINCE

	Antenatal HIV Prevalence 2007	% antenatal HIV test (%)	% admin of PMTCT to babies	Estimated Coverage (prevalence X %tested X %admin to baby)	Est. not Covered	Transmission Rate in exposed assuming SD NVP MTCT=15% & untreated MTCT=29% (Rollins)*	Overall Population Prevalence per 100 kids	Same precision across province	Same relative precision across province (%)	sample size for 30% relative precision	Sample size for design effect** of 2 & relative precision 30%	Overall population prevalence per 100 kids (%)	Varying precision by province	Varying relative precision by province (%)	Sample size using varying precision level without design effect	Sample size using varying precision level with design effect** of 2
SA	29	67	47	31.5%	68.5%	24.6%	7.1%	2.1	30	575	1150					
WC*	15	97	75	72.8%	27.3%	13.0%	1.9%	0.6	30	1989	3978	1.9	1.0	51	716	1400
NC	14	81	70	56.7%	43.3%	21.1%	2.9%	0.9	30	1336	2672	2.9	1.8	60	350	700
EC	24	73	35	25.6%	74.5%	25.4%	6.1%	1.8	30	680	1360	6.1	1.8	30	700	1400
FS	29	70	52	36.4%	63.6%	23.9%	6.9%	2.1	30	560	1120	6.9	2.0	29	617	1300
KZN	37	66	52	34.3%	65.7%	21.4%	7.9%	2.4	30	485	970	7.9	2.0	25	699	1400
MP	34	56	36	20.2%	79.8%	26.2%	8.9%	2.7	30	428	856	8.9	2.0	22	779	1600
LP	20	74	54	40.0%	60.0%	23.4%	4.7%	1.4	30	878	1756	4.7	1.5	32	703	1400
NW	29.9	86	50	43.0%	57.0%	23.0%	6.9%	2.1	30	560	1119	6.9	2.0	29	601	1200
GP	31	65	27	17.6%	82.5%	26.5%	8.2%	2.5	30	463	926	8.2	2.0	24	723	1800
Total										7379	14758					12200

ANC Prevalence & Coverage Data from DHIS

\*WC & KZN assume full coverage dual therapy - Rollins KZN Study is 7%

\*\* Design Effect =  $1+(100-1)*(ICC=.01)=2$

## SAMPLING

Table 3 - number of facilities needed to be sampled from each province to collect data within 3wks (4 weeks for Northern Cape) duration from each facility.

Note DTP1 = 1<sup>st</sup> DTP

### WESTERN CAPE

Strata	Total Annual DTPDTP#	Percentage	Sample Size Proportional	Median Yearly Clinic DTP1 Number	Media 3-week Clinic DTP1 Number	Number of Facilities to be Visited
Small clinics (< 130 DTPDTP1#)	4537					20
Medium size clinics (130-300 annual DTPDTP1#)	15953	17.85	250	192	11	23
Large size (annual DTPDTP1 #>300) but low prev clinics	62884	70.38	985	535	31	32
Large size (annual DTPDTP1 #>300) but high prev clinics	10517	11.77	165	857	49	3
Overall Total	89354	100	1400			58 (or 78 if small facilities are included)

**EASTERN CAPE**

Strata	Total Annual DTPDTP for the province	Percentage	Adjusted Percentage	Sample size proportional	Sample size adjusted proportional	Median yearly clinic DTP1 number	Median 3 week clinic DTP1 number	number of facilities need to be visited	Number of facilities that should be visited based on adjusted distribution
Small clinics (<130 DTPDTP1#)	25862							20	20
Medium size clinics (130-300 annual DTP1#)	41620	36.38	30	509	420	186.5	11	47	39
large size (Annual DTP1 #>300) but low prev clinics	41646	36.40	43	510	602	459	26	19	23
large size (Annual DTP1 #>300) but high prev clinics	31141	27	27	381	378	402	23	16	16
Overall Total	114407	100	100	1400	1400			83	78 (or 98 if small facilities are included)

**FREE STATE**

Strata	Total Annual DTP for the province	Percentage	Sample size proportional	Median yearly clinic DTP1 number	Median 3 week clinic DTP1 number	number of facilities need to be visited
Small clinics (<130 DTP1#)	4880					20
Medium size clinics (130-300 annual DTP1#)	14418	27.34%	355	201	12	31
large size (Annual DTP1 #>300) but high prev clinics	38326	72.66%	945	404	23	41
Overall Total	52744	100%	1300			72 (or 92 if small facilities are included)

**GAUTENG**

Strata	Total Annual DTP for the province	Percentage	Sample size proportional	Median yearly clinic DTP1 number	Median 3 week clinic DTP1 number	number of facilities need to be visited
Small clinics (<130 DTP1#)	1926					20
Medium size clinics (130-300 annual DTP1#)	15359	8.95%	161	237.5	14	12
Large size (Annual DTP1 #>300) but low prev clinics	33023	19.25%	347	549	32	11
Large size (Annual DTP1 #>300) but high prev clinics	123199	71.80%	1292	629	36	36
Overall Total	171581	100%	1800			59 (or 79 if small facilities are included)



**KWAZULU-NATAL**

Strata	Total Annual DTP for the province	Percentage	Sample size proportional	Median yearly clinic DTP1 number	Median 3 week clinic DTP1 number	number of facilities need to be visited
Small clinics (<130 DTP1#)	7365					20
Medium size clinics (130-300 annual DTP1#)	40070	20.84%	292	209	12	24
large size (Annual DTP1 #>300) but low prev clinics	6505	3.38%	47	536.5	31	2
large size (Annual DTP1 #>300) but high prev clinics	145661	75.77%	1061	483	28	38
Over all Total	192236	100%	1400			64 (or 84 if small facilities are included)

**LIMPOPO**

Strata	Total Annual DTP for the province	Percentage	Sample size proportional	Median yearly clinic DTP1 number	Median 3 week clinic DTP1 number	number of facilities need to be visited
Small clinics (<130 DTP1#)	7166					20
Medium size clinics (130-300 annual DTP1#)	41027	33.89%	474	206	12	40
large size (Annual DTP1 #>300) but low prev clinics	80048	66.11%	926	470.5	27	34
large size (Annual DTP1 #>300) but high prev clinics	0	0.00%	0		0	0
Over all Total	121075	100%	1400			74 (or 94 if small facilities are included)

**MPUMALANGA**

Strata	Total Annual DTP for the province	Percentage	Adjusted percentage	Sample size proportional	Sample size adjusted proportional	Median yearly clinic DTPDTP1 number	Median 3 week clinic DTPDTP1 number	number of facilities need to be visited	number of facilities need to be visited based on adjusted distribution
Small clinics (<130 DTP1#)	4545							20	
Medium size clinics (130-300 annual DTP1#)	20858	26.73%	20%	428	320	225	13	33	25
large size (Annual DTP1 #>300) but low prev clinics	0	0.00%		0	0	0	0	0	0
large size (Annual DTP1 #>300) but high prev clinics	57172	73.27%	80%	1172	1280	439	25	46	51
Overall Total	78030	100%	100%	1600	1600			79	76

**NORTHERN CAPE**

Strata	Total Annual DTP for the province	Percentage	Sample size proportional	Median yearly clinic DTP1 number	Median 4 week clinic DTP1 number	number of facilities need to be visited
Small clinics (<130 DTP1#)	2475					20
Medium size clinics (130-300 annual DTP1#)	7766	51.82%	363	207.5	16	23
large size (Annual DTP1 #>300) but low prev clinics	7221	48.18%	337	400	32	11
large size (Annual DTP1 #>300) but high prev clinics	0	0.00%	0		0	
Overall Total	14987	100%	700			34 (or 54 if small facilities are included)

**NORTH WEST**

Strata	Total Annual DTP for the province	Percentage	Sample size proportional	Median yearly clinic DTP1 number	Median 3 week clinic DTP1 number	number of facilities need to be visited
Small clinics (<130 DTP1#)	8758					20
Medium size clinics (130-300 annual DTP1#)	22925	34.26%	411	204.5	12	35
large size (Annual DTP1 #>300) but low prev clinics	24100	36.02%	432	413	24	18
large size (Annual DTP1 #>300) but high prev clinics	19887	29.72%	357	432.5	25	14
Over all Total	66912	100%	1200			67 (or 87 if small facilities are included)

Each province was divided into three strata:

- Stratum 1 includes clinics and CHCs that have annual Dtp1st dose 130-300 based on the 2007 DHIS data.
- Stratum 2 includes clinics & CHCs with 300 and above Dtp1st dose & HIV prevalence below the national (<29%) rate based on the 2007 DHIS and 2008 antenatal survey data respectively.
- Stratum 3 includes clinics & CHCs with 300 & above Dtp1st dose (based on the 2007 DHIS data) & HIV prevalence above the national rate based on the 2007 DHIS data & 2008 antenatal survey data respectively.

In the small facilities stratum, 20 facilities were selected for situational analysis. Out of these 20 facilities only 10 were selected based on the feedback from the situational analysis.

#### *Provinces that do not have the third stratum*

Western Cape, Limpopo and Northern Cape do not have a third stratum: there is no district with >29% HIV prevalence and high delivery rate (>300 immunisation) in these provinces. However, for Western Cape, we had a sub-district level data from the ANC survey, which indicated that Khayelitsha sub-district has more than 29% HIV prevalence. So, the third stratum was created from large clinics in Khayelitsha. We were not able to do the same for Limpopo and Northern Cape, as we did not have sub-district level HIV prevalence data (from the ANC survey) for the two provinces.

#### *Reduced sample size for Northern Cape to 700*

There were 96 facilities that could be sampled in the Northern Cape; but logistically 96 clinics was not an achievable target within the allocated time. Consequently the researchers reduced the number of facilities that needed to be visited to 53.

#### *Adjusting weighting for Mpumalanga and Eastern Cape*

The number of facilities needed to be sampled for Mpumalanga and Eastern Cape was 79 and 83 respectively. In addition, most of the facilities were from medium sized clinics. As this might be difficult to achieve with available logistics capacity, we slightly shifted the weighting to the large clinics (see column D), hence the number of facilities needed to be sampled from medium facilities decreased from 47 to 39 for the Eastern Cape and from 33 to 25 for Mpumalanga (see column J). This was logistically feasible.

In Free state, we had only two strata - we grouped the last two strata as one stratum.

The second strata (large and low HIV prevalence clinics) in Free state had only 0.74% weighting which translates to sampling only 1 facility from the second stratum. Since sampling cannot be done for one facility, the second stratum is combined with the third stratum thus we have only two strata for Free state.

APPENDIX 2

wc Wellington CDC						
wc Dr Ivan Toms						
wc Bongolethu Clinic	Public	256	15	1	11	* note: have removed Kwanokut hula clinic

mp Boekenhouthoek Clinic	Public	174	10	1	13	Wesselton Clinic removed
mp Gemsbokspruit Clinic	Public	136	8	1	13	Kriel Clinic removed replace with????

gp Joy Clinic		290	17	14		this is replacement for gp Bapsfontein Clinic
---------------	--	-----	----	----	--	--

ec Oliver Tambo District Municipality	ec Nyandeni Local Service Area	ec Nolita Clinic	Public	153
---------------------------------------	--------------------------------	------------------	--------	-----

ec Chris Hani District Municipality	ec Ngcobo Local Service Area	ec Lahlangubo Clinic (Ngcobo)	Public	279
-------------------------------------	------------------------------	-------------------------------	--------	-----

kz Umkhanyakude District Municipality	kz KwaMbuzi Clinic	Public	254
kz Umkhanyakude District Municipality	kz Ophondweni Clinic	Public	167