

REPORT ON WEEKLY DEATHS IN SOUTH AFRICA

16 – 22 JAN 2022

(WEEK 3)

Debbie Bradshaw, Ria Laubscher,
Rob Dorrington, Pam Groenewald, Tom Moultrie

Burden of Disease Research Unit
South African Medical Research Council
25 January 2022



UCT Centre
for Actuarial
Research

Glossary:

Age-standardised excess death rate: Indirectly age-standardised excess death rates have been calculated for each province to adjust the crude death rates per capita for the differences in distribution of the population by age. The adjustment factor for each province is calculated as the crude death rate for South Africa divided by what the crude rate for South Africa would have been had the age distribution of the population been that of the province. Standardisation for age is necessary when comparing populations that differ in their age structure because age has a powerful influence on the risk of dying. The rate is based on the cumulative number of excess deaths since 3 May 2020 to date divided by the population estimate for 2021 and has not been annualised.

Actual number of deaths: The actual number of deaths in South Africa have been estimated from the numbers recorded on the National Population Register using weighting factors set to produce results consistent with those of the annual Rapid Mortality Surveillance Report to account for deaths of persons who are not on the National Population Register as well as those that have not been registered with the Department of Home Affairs. **The adjustments to account for incompleteness of recording of deaths on the NPR have been re-estimated for the 2021 reports taking into account the 2017 cause-of-death data released by Stats SA in 2020.** A methodological note briefly outlining the changes can be downloaded with this report from the SAMRC website: <https://www.samrc.ac.za/reports/report-weekly-deaths-south-africa>.

Epi-week: The Weekly Death Reports in 2020 used weeks from 1 January and ran from Wednesday to Tuesday. In setting up the monitoring for 2021, we recast the data to report by an 'Epi-week' consistent with CDC and many NICD reports which run from Sunday to Saturday, ensuring continuity of weeks from one year to the next. Each week is aligned with the 'Epi-year' that has 4 or more days in that week. Week 53 of 2020 is from 27 December 2020 to 2 January 2021, Week 1 of 2021 is 3 January – 9 January 2021 and Week 1 of 2022 is 2 January – 8 January 2022.

Excess deaths: There is no universal definition of, or understanding of what is meant by, "excess mortality". It is a term used in epidemiology and public health that refers to the number of deaths that are occurring above what we would normally expect. The WHO uses the term to describe "Mortality above what would be expected based on the non-crisis mortality rate in the population of interest. Excess mortality is thus mortality that is attributable to the crisis conditions. It can be expressed as a rate (the difference between observed and non-crisis mortality rates), or as a total number of excess deaths."

Excess natural deaths associated with COVID-19: Generally, the number of excess deaths per week is calculated as the number of all-cause deaths in that week less the number that might be assumed to have occurred had there not been the epidemic (i.e. the counterfactual number), provided that the counterfactual is lower. However, this approach has generally only been applied to countries where deaths have been tracking the counterfactual before the onset of significant numbers of COVID-19 related deaths. The method provides a poor estimate of the numbers of COVID-19 and collateral deaths in the early stages of the epidemic when this is not the case. Thus, we estimated the numbers of COVID-19 and collateral deaths, once a clear upward trend is evident, as the number of actual deaths less a baseline number determined as a proportion of the predicted number. By the end of the 1st wave of the pandemic, the predicted values have been used as the counterfactual.

Warning: The Department of Home Affairs has faced sporadic temporary office closures, particularly in areas that are more affected by COVID-19. This may affect our allocation of deaths to a particular metro area. For example, a death that occurred in the City of Cape Town might have been registered at an office outside of the City because of a temporary closure of offices in the metro. Closure may also cause a delay in the processing of the death registration which would result in an underestimate of the deaths in the most recent week.

Background

This report provides estimates of the weekly number of deaths of all persons in South Africa for epidemiological **Week 3** of 2022, covering the period **16 – 22 Jan 2022**.

The predicted numbers of weekly deaths in 2022 have been estimated using the models prepared for 2021 with population estimates for 2022. A methodological note outlining the approach for monitoring deaths during 2021 can be downloaded from the SAMRC website: [https://www.samrc.ac.za/sites/default/files/files/2021-01-24/Methodological Note on Predicted Weekly Deaths 20 Jan 2021.pdf](https://www.samrc.ac.za/sites/default/files/files/2021-01-24/Methodological%20Note%20on%20Predicted%20Weekly%20Deaths%20Jan%202021.pdf).

Briefly, the predicted values for the provinces and nationally are based on negative binomial models based on death data for the period 2014-2019. After reviewing trends in the data, separate negative binomial models have been fitted to the unnatural deaths for all provinces combined, and separate models for natural deaths for each of KwaZulu-Natal, Western Cape and the Northern Cape, and the 6 other provinces in a combined model to provide estimates by age, sex and epi-week for each year. A prediction interval has been estimated on the basis of the variability in the observed weekly data for each reported domain. Predicted values for each metropolitan area continue to be based on the time-series model of data for 2018 and 2019 developed for 2020 as the trends in the sub-provincial data need further investigation to develop a comprehensive district-level model.

Trends

- The weekly number of deaths (all ages) from all causes decreased to **10,640** in **Week 3** (16 – 22 Jan 2022), resulting in an excess of **1,940** deaths during the week.
- The number of excess deaths from natural causes (all ages) peaked in **Week 51** of 2021 (19 Dec – 25 Dec 2021) and decreased to **2,657** in **Week 1** of 2022 (2 – 8 Jan 2022), and further decreased to 1,726 in **Week 3** (16 – 22 Jan 2022) with a p-score of 22% (i.e., the excess was 22% of the predicted number).
- Since 3 May 2020, there has been a cumulative total of more than **294,000** excess deaths from natural causes of persons all ages of which **85,000** occurred in 2020, just over **202,000** occurred in 2021 (since 3 Jan 2021) and more than **6,000** occurred in 2022.


Week	Date	Weekly excess deaths from natural causes (all ages)	Cumulative excess since 3 May 2020 (all ages)	Cumulative excess since 3 January 2021 (all ages)	Cumulative excess since 2 January 2022 (all ages)
43	24-Oct-21 – 30-Oct-21	823	268,963	184,020	
44	31-Oct-21 – 6-Nov-21	1,414	270,377	185,433	
45	7-Nov-21 – 13-Nov-21	1,537	271,914	186,971	
46	14-Nov-21 – 20-Nov-21	1,243	273,157	188,214	
47	21-Nov-21 – 27-Nov-21	1,205	274,362	189,419	
48	28-Nov-21 – 4-Dec-21	1,880	276,242	191,299	
49	5-Dec-21 – 11-Dec-21	1,701	277,944	193,000	
50	12-Dec-21 – 18-Dec-21	2,574	280,518	195,574	
51	19-Dec-21 – 25-Dec-21	3,513	284,031	199,087	
52	26-Dec-21 – 1-Jan-22	3,510	287,541	202,598	
1	2-Jan-22 – 8-Jan-22	2,657	290,198		2,657
2	9-Jan-22 – 15-Jan-22	2,154	292,352		4,798
3	16-Jan-22 – 22-Jan-22	1,726	294,079		6,126

- For people under the age of 60, the number of natural deaths was within the prediction bounds in **Week 3** (16 – 22 Jan 2022), following an erratic peaking during the 4th wave. cumulative number of excess natural deaths for people under-60 years since **3 May 2020** was just over **77,000**.
- The number of weekly excess deaths in the 60+ years age group increased to a peak of **7,160** in **Week 29** (18 – 24 Jul 2021) in wave 3 with a p-score of 154%. After an erratic decline, the numbers remained above the upper prediction bound but reached a low in **Week 43** (24 – 30 Oct 2021) with a p-score of 17%. The weekly number of excess deaths in the age group increased during Wave 4 and have decreased to about **1,328** with a p-score of about 35% in **Week 3** (16 – 22 Jan 2022). The cumulative total number of excess natural deaths in this age group since **3 May 2020** was just under **217,000**.

Week	Date	Weekly excess deaths from natural causes for persons 60+ years	p-score
43	24-Oct-21 – 30-Oct-21	697	16.8%
44	31-Oct-21 – 6-Nov-21	838	20.9%
45	7-Nov-21 – 13-Nov-21	937	23.6%
46	14-Nov-21 – 20-Nov-21	844	21.8%
47	21-Nov-21 – 27-Nov-21	814	21.2%
48	28-Nov-21 – 4-Dec-21	1,116	28.0%
49	5-Dec-21 – 11-Dec-21	1,201	29.7%
50	12-Dec-21 – 18-Dec-21	1,757	45.6%
51	19-Dec-21 – 25-Dec-21	2,437	60.8%
52	26-Dec-21 – 1-Jan-22	2,586	63.7%
1	2-Jan-22 – 8-Jan-22	2,140	51.1%
2	9-Jan-22 – 15-Jan-22	1,654	42.3%
3	16-Jan-22 – 22-Jan-22	1,328	34.6%

- The 4th wave has peaked in all of the nine provinces and numbers of natural deaths have decreased with small fluctuations. However, deaths from natural causes in **Week 3** (16 – 22 Jan 2022) were higher than their upper prediction bound for all provinces except **Gauteng** and **North West**.
- Deaths from natural causes in **Buffalo City, Cape Town, Mangaung** and **Nelson Mandela Bay** were still higher than their upper prediction bound in **Week 3** (16 – 22 Jan 2022).
- Per capita excess death rates have been calculated for the provinces to scale the cumulative deaths for the population size of each province (**Table 1**). By the end of **Week 3** (16 – 22 Jan 2022), the national excess death rate since 3 May 2020 was **494** per **100,000** population.
- The provinces with the highest cumulative numbers of excess deaths at the end of **Week 3** (16 – 22 Jan 2022), are, in order, **KwaZulu-Natal, Gauteng** and **Eastern Cape**. The ranking changes to **Eastern Cape, Northern Cape** and **Free State** for the crude death rates per capita (i.e., taking size of the provincial populations into account) and to **Northern Cape** and **KwaZulu-Natal/Eastern Cape** using the age-standardised rates per capita (i.e., taking into account the age distribution of the provincial population). The **Western Cape** followed by **Gauteng** continue to have the lowest cumulative age standardised per capita rates.
- Following some declines below the predicted, coinciding with the change of lockdown to adjusted level 4 with re-banning of alcohol sales and extension of curfew, the number of unnatural deaths has tracked the

predicted number since **Week 30** (25 – 31 Jul 2021). **Week 52** (26 Dec 2021 – 1 Jan 2022), followed the predicted trend closely and saw the highest number of unnatural deaths in 2021, at **1,574**. In **Week 3** (16 – 22 Jan 2022), there were **1,115** unnatural deaths, higher than the predicted number.



Numbers have been scaled to the estimated actual number of death and for the last week has been adjusted for delayed registrations




Table 1: Number of excess natural deaths of persons by province and metro relative to revised predicted number based on the observed drop during lockdown, South Africa 2020/21

Region	Period	Excess deaths vs revised base	Excess deaths per 100,000 population	Age standardised excess death rate per 100,000
South Africa	3 May 20 – 22 Jan 22	294,079	494	494
Province				
Eastern Cape	31 May 20 – 22 Jan 22	49,745	756	609
Free State	21 Jun 20 – 22 Jan 22	16,559	569	569
Gauteng	7 Jun 20 – 22 Jan 22	58,134	373	410
KwaZulu-Natal	7 Jun 20 – 22 Jan 22	60,505	529	608
Limpopo	21 Jun 20 – 22 Jan 22	31,327	530	464
Mpumalanga	21 Jun 20 – 22 Jan 22	22,628	470	507
Northern Cape	28 Jun 20 – 22 Jan 22	8,344	713	667
North West	28 Jun 20 – 22 Jan 22	16,556	411	422
Western Cape	3 May 20 – 22 Jan 22	30,282	429	378
Metropolitan Municipality				
Buffalo City	31 May 20 – 22 Jan 22	5,572		
City of Cape Town	3 May 20 – 22 Jan 22	21,593		
Ekurhuleni	7 Jun 20 – 22 Jan 22	14,380		
eThekweni	14 Jun 20 – 22 Jan 22	13,374		
Johannesburg	7 Jun 20 – 22 Jan 22	19,876		
Mangaung	21 Jun 20 – 22 Jan 22	4,847		
Nelson Mandela Bay	31 May 20 – 22 Jan 22	7,711		
City of Tshwane	7 Jun 20 – 22 Jan 22	11,020		

Note: Period has been determined based on when an upturn in the number of natural deaths became apparent. Parts do not sum to the whole because office closures due to Covid-19 may have led to registration of deaths at other offices which may not be in the same area, and random fluctuation at the point at which the baseline is determined.


Table 2: Number of excess deaths from all causes of persons by province and metro relative to predicted number based on historical trend, South Africa 2020/21

Region	Excess deaths vs forecast	Excess deaths per 100,000 population
South Africa	284,556	478
Province		
Eastern Cape	49,123	746
Free State	15,885	546
Gauteng	54,916	352
KwaZulu-Natal	60,318	527
Limpopo	30,769	521
Mpumalanga	21,933	456
Northern Cape	8,051	688
North West	15,804	392
Western Cape	27,756	393
Metropolitan Municipality		
Buffalo City	5,443	
City of Cape Town	19,454	
Ekurhuleni	13,082	
eThekweni	12,091	
Johannesburg	19,444	
Mangaung	5,070	
Nelson Mandela Bay	7,459	
City of Tshwane	11,238	




Numbers have been scaled to the estimated actual number of death and for the last week has been adjusted for delayed registrations


Eastern Cape weekly deaths from natural causes : 29 Dec 2019 - 22 Jan 2022






Free State weekly deaths from natural causes : 29 Dec 2019 - 22 Jan 2022






Gauteng weekly deaths from natural causes : 29 Dec 2019 - 22 Jan 2022






Numbers have been scaled to the estimated actual number of death and for the last week has been adjusted for delayed registrations






Numbers have been scaled to the estimated actual number of death and for the last week has been adjusted for delayed registrations





Numbers have been scaled to the estimated actual number of death and for the last week has been adjusted for delayed registrations




Numbers have been scaled to the estimated actual number of death and for the last week has been adjusted for delayed registrations



Numbers have been scaled to the estimated actual number of death and for the last week has been adjusted for delayed registrations



Numbers have been scaled to the estimated actual number of death and for the last week has been adjusted for delayed registrations



Numbers have been scaled to the estimated actual number of death and for the last week has been adjusted for delayed registrations. As only a quarter to a third of unnatural deaths in the most recent week are processed at the time of the survey, the estimate for the most recent week is quite uncertain.

Vertical lines in order

- 0 Week Disaster Management Act implemented
- 1 Week lockdown level 5 introduced
- 2 Week lockdown changed to level 4, with curfew
- 3 Week lockdown changed to level 3 including unbanning of alcohol
- 4 Week alcohol re-banned and a curfew re-introduced
- 5 Week lockdown changed to level 2, including unbanning of alcohol
- 6 Week lockdown changed to level 1
- 7 Week lockdown changed to level 3 advanced (rebanning alcohol and a extension of curfew)
- 8 Week lockdown relaxed to allow sale of alcohol 4 days/week and reduce curfew
- 9 Week lockdown relaxed to allow sale of alcohol except during curfew and reduce curfew to midnight to 4am
- 10 Week lockdown changed to level 3 advanced (limiting alcohol and a extending of curfew)
- 11 Week lockdown changed to level 4, with re-banning of alcohol, curfew 9pm-4am
- 12 Week of unrest in KZN and GT
- 13 Week lockdown changed to level 3 advanced (alcohol 4 days/w, curfew 10pm-4am)
- 14 Week lockdown changed to level 2 advanced (alcohol 5 days/w, curfew 11pm-4am)
- 15 Week lockdown changed to level 1 advanced (no alcohol post 11pm, curfew 12pm-4am, large gatherings)
- 16 Week lockdown level 1 advanced (removed limits on alcohol & curfew, allowed larger gatherings)